

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 June 2003 (26.06.2003)

PCT

(10) International Publication Number
WO 03/051905 A2

(51) International Patent Classification⁷: C07K

(74) Agents: SHAYESTEH, Laleh et al.; Exelixis, Inc., P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(21) International Application Number: PCT/US02/39742

(22) International Filing Date:
12 December 2002 (12.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/340,312 13 December 2001 (13.12.2001) US

(71) Applicant: EXELIXIS, INC. [US/US]; P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(72) Inventors: COSTA, Michael, A.; 18 Hazelwood Avenue, San Francisco, CA 94112 (US). GENDREAU, Steven, Brian; 2801 Turk Street, #103, San Francisco, CA 94118 (US). DORA, Emery, G., III; 1847 32nd Avenue, San Francisco, CA 94122 (US). NICOLL, Monique; 224 Naomi Avenue, Pacifica, CA 94044 (US). URBANI, Lenore; 59 Playbowl Drive, La Honda, CA 94020 (US). LARSON, Jeffrey, S.; 1220 El Camino Real #305, Burlingame, CA 94010 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TAOJKS AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

(57) Abstract: Human TAOJK genes are identified as modulators of the beta-catenin pathway, and thus are therapeutic targets for disorders associated with defective beta-catenin function. Methods for identifying modulators of beta-catenin, comprising screening for agents that modulate the activity of TAOJK are provided.

WO 03/051905 A2

TAOJIKs AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to U.S. provisional patent application 60/340,312 filed 12/13/2001. The content of the prior application is hereby incorporated in its entirety.

BACKGROUND OF THE INVENTION

10 Beta-catenin is an adherens junction protein. Adherens junctions (AJs; also called the zonula adherens) are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces. AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior. At several stages of
15 embryogenesis, wound healing, and tumor cell metastasis, cells form and leave epithelia. This process, which involves the disruption and reestablishment of epithelial cell-cell contacts, may be regulated by the disassembly and assembly of AJs. AJs may also function in the transmission of the 'contact inhibition' signal, which instructs cells to stop dividing once an epithelial sheet is complete.

20 The AJ is a multiprotein complex assembled around calcium-regulated cell adhesion molecules called cadherins (Peifer, M.(1993) Science 262: 1667-1668). Cadherins are transmembrane proteins: the extracellular domain mediates homotypic adhesion with cadherins on neighboring cells, and the intracellular domain interacts with cytoplasmic proteins that transmit the adhesion signal and anchor the AJ to the actin
25 cytoskeleton. These cytoplasmic proteins include the alpha-, beta-, and gamma-catenins. The beta-catenin protein shares 70% amino acid identity with both plakoglobin, which is found in desmosomes (another type of intracellular junction), and the product of the Drosophila segment polarity gene 'armadillo'. Armadillo is part of a multiprotein AJ complex in Drosophila that also includes some homologs of alpha-catenin and cadherin,
30 and genetic studies indicate that it is required for cell adhesion and cytoskeletal integrity.

Beta-catenin, in addition to its role as a cell adhesion component, also functions as a transcriptional co-activator in the Wnt signaling pathway through its interactions with the family of Tcf and Lef transcription factors (for a review see Polakis, (1999) Current Opinion in Genetics & Development, 9:15-21 and Gat U., et al., (1998) Cell 95:605-614).

The APC gene, which is mutant in adenomatous polyposis of the colon, is a negative regulator of beta-catenin signaling (Korinek, V. et al., (1997) Science 275: 1784-1787; Morin, P. J., et al., (1997) Science 275: 1787-1790). The APC protein normally binds to beta-catenin and, in combination with other proteins (including glycogen synthase kinase-3b and axin, is required for the efficient degradation of b-catenin. The regulation of beta-catenin is critical to the tumor suppressive effect of APC and that this regulation can be circumvented by mutations in either APC or beta-catenin.

While mammals contain only a single beta-catenin gene, *C. elegans* contains three (Korswagen HC, et al., (2000) Nature 406:527-32). Each worm beta-catenin appears to carry out unique functions (Korswagen HC, et al., (2000) Nature 406:527-32, Nartarajan L et al. (2001) Genetics 159: 159-72). Because of the divergence of function in *C. elegans*, it is possible to specifically study beta-catenin role in cell adhesion, which is mediated by the *C. elegans* beta-catenin HMP-2.

Eukaryotic cells respond to extracellular stimuli by recruiting signal transduction pathways, many of which employ protein Ser/Thr kinases of the ERK family (Levin, D. E., and Errede, B. (1995) Curr. Opin. Cell Biol. 7, 197-202). The ubiquity of ERKs and their upstream activators, the MEKs, in signal transduction was first appreciated from studies of yeast (Herskowitz, I. (1995) Cell 80, 187-197). Part of the cellular response to toxins, physical stresses and inflammatory cytokines occurs by signalling via the stress-activated protein kinase (SAPK) and p38 reactivating kinase pathways (Kyriakis, J. M., and Avruch, J. (1990) J. Biol. Chem. 265, 17355-17363; Kyriakis, J. M., et al., (1991) J. Biol. Chem. 266, 10043-10046; Pulverer, B. J., et al., (1991) Nature 353, 670-674). These stress-responsive kinase pathways are structurally similar, but functionally distinct, from the archetypal mitogen-activated protein kinases (MAPKs or ERKs). The stimuli that start the pathway result in modification of cellular gene expression, growth arrest, apoptosis, or activation of immune and reticuloendothelial cells. TAO1 (thousand and one amino acid) is a protein kinase that may play a role in regulating stress-responsive MAP kinase pathways (Hutchison, M., et al (1998). J Biol Chem 273:28625-32). KIAA1361 is a protein with strong similarity with TAO1. JIK (JNK-SAPK inhibitory kinase) is an STE20-like serine/threonine kinase and member of the GCK-like subfamily of Ste20 kinases. JIK is activated by ligand-bound EGF receptors, inhibits the JNK/SAPK signaling pathway, and also interacts with IRE1 (a gene involved in endoplasmic reticulum stress response) (Tassi, E., et al (1999) J Biol Chem 274:33287-95; Zhang, W.,

et al (2000) Biochem Biophys Res Commun 274:872-9; Yoneda, T., et al (2001) J Biol Chem 276:13935-40).

The ability to manipulate the genomes of model organisms such as *C. elegans* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Dulubova I, et al, J Neurochem 2001 Apr;77(1):229-38; Cai T, et al., Diabetologia 2001 Jan;44(1):81-8; Pasquinelli AE, et al., Nature. 2000 Nov 2;408(6808):37-8; Ivanov IP, et al., EMBO J 2000 Apr 17;19(8):1907-17; Vajo Z et al., Mamm Genome 1999 Oct;10(10):1000-4). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a "genetic entry point") that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a "modifier" involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as beta-catenin, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

SUMMARY OF THE INVENTION

We have discovered genes that modify the beta-catenin pathway in *C. elegans*, and identified their human orthologs, hereinafter referred to as TAO and JIK related kinases (TAOJIK). The invention provides methods for utilizing these beta-catenin modifier genes and polypeptides to identify TAOJIK-modulating agents that are candidate therapeutic agents that can be used in the treatment of disorders associated with defective or impaired beta-catenin function and/or TAOJIK function. Preferred TAOJIK-modulating agents specifically bind to TAOJIK polypeptides and restore beta-catenin

function. Other preferred TAOJIK-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress TAOJIK gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

5 TAOJIK modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with a TAOJIK polypeptide or nucleic acid. In one embodiment, candidate TAOJIK modulating agents are tested with an assay system comprising a TAOJIK polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate beta-catenin
10 modulating agents. The assay system may be cell-based or cell-free. TAOJIK-modulating agents include TAOJIK related proteins (e.g. dominant negative mutants, and biotherapeutics); TAOJIK-specific antibodies; TAOJIK-specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind to or interact with TAOJIK or compete with TAOJIK binding partner (e.g. by binding to a TAOJIK
15 binding partner). In one specific embodiment, a small molecule modulator is identified using a kinase assay. In specific embodiments, the screening assay system is selected from a binding assay, an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

In another embodiment, candidate beta-catenin pathway modulating agents are
20 further tested using a second assay system that detects changes in the beta-catenin pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals
25 predetermined to have a disease or disorder implicating the beta-catenin pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

The invention further provides methods for modulating the TAOJIK function and/or the beta-catenin pathway in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a TAOJIK polypeptide or nucleic acid. The agent
30 may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated the beta-catenin pathway.

DETAILED DESCRIPTION OF THE INVENTION

Genetic screens were designed to identify modifiers of the beta-catenin pathway in *C. elegans*. A weak allele of beta-catenin was used in our screen (a homozygous viable mutant of beta-catenin, allele qm39). The hmp-2 (qm-39) strain produces larval worms with a highly penetrant lumpy body phenotype in first stage larval worms (L1s). Various specific genes were silenced by RNA inhibition (RNAi). Methods for using RNAi to silence genes in *C. elegans* are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); WO9932619). Genes causing altered phenotypes in the worms were identified as modifiers of the beta-catenin pathway. A modifier of particular interest was T17E9.1. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, TAOJIK genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective beta-catenin signaling pathway, such as cancer.

In vitro and in vivo methods of assessing TAOJIK function are provided herein. Modulation of the TAOJIK or their respective binding partners is useful for understanding the association of the beta-catenin pathway and its members in normal and disease conditions and for developing diagnostics and therapeutic modalities for beta-catenin related pathologies. TAOJIK-modulating agents that act by inhibiting or enhancing TAOJIK expression, directly or indirectly, for example, by affecting a TAOJIK function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. TAOJIK modulating agents are useful in diagnosis, therapy and pharmaceutical development.

Nucleic acids and polypeptides of the invention

Sequences related to TAOJIK nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) number) as GI#s 4759207 (SEQ ID NO:1), 15929548 (SEQ ID NO:2), 7706400 (SEQ ID NO:4), 7705559 (SEQ ID NO:5), 20552662 (SEQ ID NO:6), 12803830 (SEQ ID NO:7), 19923463 (SEQ ID NO:8), 15302531 (SEQ ID NO:9), 7243102 (SEQ ID NO:10), 18587661 (SEQ ID NO:11), 20559660 (SEQ ID NO:12), 11596143 (SEQ ID NO:13), and 12803830 (SEQ ID NO:15) for nucleic acid, and GI#s 4759208 (SEQ ID NO:16), 7705560 (SEQ ID NO:17), and 7243103 (SEQ ID NO:18) for polypeptides. Additionally, nucleic acid sequences of SEQ ID NOs:3 and 14 can also be used in the invention.

TAOJIKs are kinase proteins with kinase domains. The term "TAOJIK polypeptide" refers to a full-length TAOJIK protein or a functionally active fragment or derivative thereof. A "functionally active" TAOJIK fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type TAOJIK protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of TAOJIK proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one embodiment, a functionally active TAOJIK polypeptide is a TAOJIK derivative capable of rescuing defective endogenous TAOJIK activity, such as in cell based or animal assays; the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of a TAOJIK, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., *et al.*, Nucleic Acids Res, 1999, 27:260-2). For example, the kinase (PFAM 00069) domain of TAOJIK from GI#s 4759208, 7705560, and 7243103 (SEQ ID NOs:16, 17, and 18, respectively) are located at approximately amino acid residues 28 to 281, 24 to 277, and 32 to 285, respectively. Methods for obtaining TAOJIK polypeptides are also further described below. In some embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of any one of SEQ ID NOs:16-18 (a TAOJIK). In further preferred embodiments, the fragment comprises the entire kinase (functionally active) domain.

The term "TAOJIK nucleic acid" refers to a DNA or RNA molecule that encodes a TAOJIK polypeptide. Preferably, the TAOJIK polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with human TAOJIK. Methods of identifying orthologs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result

retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD *et al.*, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight
5 conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by
10 ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *C. elegans*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion
15 of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all
20 the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which
25 the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted
30 for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and

histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; database: European Bioinformatics Institute; Smith and Waterman, 1981, *J. of Molec. Biol.*, 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov, 1986 *Nucl. Acids Res.* 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of any of SEQ ID NOs:1-15. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are set out in readily available procedure texts (*e.g.*, *Current Protocol in Molecular Biology*, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of any one of SEQ ID NOs:1-15 under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5mM EDTA, 0.1% PVP, 0.1%

Ficoll, 1% BSA, and 500 μ g/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 μ g/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

Isolation, Production, Expression, and Mis-expression of TAOJIK Nucleic Acids and Polypeptides

TAOJIK nucleic acids and polypeptides, useful for identifying and testing agents that modulate TAOJIK function and for other applications related to the involvement of TAOJIK in the beta-catenin pathway. TAOJIK nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of a TAOJIK protein for assays used to assess TAOJIK function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) Protein Expression: A Practical Approach, Oxford University Press Inc., New York 1999; Stanbury PF et al., Principles of Fermentation Technology, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) Protein Purification Protocols, Humana Press, New Jersey, 1996; Coligan JE et al, Current Protocols in Protein Science (eds.), 1999, John Wiley & Sons, New

York). In particular embodiments, recombinant TAOJIK is expressed in a cell line known to have defective beta-catenin function. The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

5 The nucleotide sequence encoding a TAOJIK polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native TAOJIK gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus);
10 microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the TAOJIK gene product, the expression vector can comprise a promoter operably linked to a TAOJIK gene nucleic acid, one or more origins
15 of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the TAOJIK gene product based on the physical or functional properties of the TAOJIK protein in *in vitro* assay systems (*e.g.* immunoassays).

20 The TAOJIK protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and
25 expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the TAOJIK gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange,
30 affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native TAOJIK proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as

immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of TAOJK or other genes associated with the beta-catenin pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

Genetically modified animals

Animal models that have been genetically modified to alter TAOJK expression may be used in *in vivo* assays to test for activity of a candidate beta-catenin modulating agent, or to further assess the role of TAOJK in a beta-catenin pathway process such as apoptosis or cell proliferation. Preferably, the altered TAOJK expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal TAOJK expression. The genetically modified animal may additionally have altered beta-catenin expression (*e.g.* beta-catenin knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, *i.e.* mosaic animals (see, for example, techniques described by Jakobovits, 1994, *Curr. Biol.* 4:761-763.) or stably integrated into its germ line DNA (*i.e.*, in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster *et al.*, *Proc. Nat. Acad. Sci. USA* 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Pat. No. 4,873,191 by Wagner *et al.*, and Hogan, B., *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, *Science* (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) *Nature* 402:370-371; for transgenic

Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous TAOJIK gene that results in a decrease of TAOJIK function, preferably such that TAOJIK expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse TAOJIK gene is used to construct a homologous recombination vector suitable for altering an endogenous TAOJIK gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, Science (1989) 244:1288-1292; Joyner *et al.*, Nature (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, Science (1989) 244:1281-1288; Simms *et al.*, Bio/Technology (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) Scan J Immunol 40:257-264; Declerck PJ *et al.*, (1995) J Biol Chem 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the TAOJIK gene, e.g., by introduction of additional copies of TAOJIK, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the TAOJIK gene. Such regulatory

sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that
5 may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two
10 transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the
15 transgene, and for sequential deletion of vector sequences in the same cell (Sun X *et al.* (2000) Nat Genet 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the beta-catenin pathway, as animal models of disease and disorders implicating defective beta-catenin function, and for *in vivo* testing of candidate therapeutic agents, such as those
20 identified in screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered TAOJIK function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered TAOJIK expression that receive candidate therapeutic agent.

25 In addition to the above-described genetically modified animals having altered TAOJIK function, animal models having defective beta-catenin function (and otherwise normal TAOJIK function), can be used in the methods of the present invention. For example, a beta-catenin knockout mouse can be used to assess, *in vivo*, the activity of a candidate beta-catenin modulating agent identified in one of the *in vitro* assays described
30 below. Preferably, the candidate beta-catenin modulating agent when administered to a model system with cells defective in beta-catenin function, produces a detectable phenotypic change in the model system indicating that the beta-catenin function is restored, i.e., the cells exhibit normal cell cycle progression.

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate the function of TAOJIK and/or the beta-catenin pathway. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the beta-catenin pathway, as well as in further analysis of the TAOJIK protein and its contribution to the beta-catenin pathway. Accordingly, the invention also provides methods for modulating the beta-catenin pathway comprising the step of specifically modulating TAOJIK activity by administering a TAOJIK-interacting or -modulating agent.

As used herein, a "TAOJIK-modulating agent" is any agent that modulates TAOJIK function, for example, an agent that interacts with TAOJIK to inhibit or enhance TAOJIK activity or otherwise affect normal TAOJIK function. TAOJIK function can be affected at any level, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a preferred embodiment, the TAOJIK - modulating agent specifically modulates the function of the TAOJIK. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the TAOJIK polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the TAOJIK. These phrases also encompasses modulating agents that alter the interaction of the TAOJIK with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of a TAOJIK, or to a protein/binding partner complex, and altering TAOJIK function). In a further preferred embodiment, the TAOJIK- modulating agent is a modulator of the beta-catenin pathway (e.g. it restores and/or upregulates beta-catenin function) and thus is also a beta-catenin-modulating agent.

Preferred TAOJIK-modulating agents include small molecule compounds; TAOJIK-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

Small molecule modulators

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight less than 10,000, preferably less than 5,000, more preferably less than 1,000, and most preferably less than 500. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the TAOJIK protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for TAOJIK-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the beta-catenin pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

Protein Modulators

Specific TAOJIK-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the beta-catenin pathway and related disorders, as well as in validation assays for other TAOJIK-modulating agents. In a preferred embodiment, TAOJIK-interacting proteins affect normal TAOJIK function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, TAOJIK-interacting proteins are useful in detecting and providing information about the function of TAOJIK proteins, as is relevant to beta-catenin related disorders, such as cancer (e.g., for diagnostic means).

An TAOJIK-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with a TAOJIK, such as a member of the TAOJIK pathway that modulates TAOJIK expression, localization, and/or activity. TAOJIK-modulators include dominant negative forms of TAOJIK-interacting proteins and of TAOJIK proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous TAOJIK-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3rd, Trends Genet (2000) 16:5-8).

An TAOJIK-interacting protein may be an exogenous-protein, such as a TAOJIK-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory; Harlow and Lane (1999) Using antibodies: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). TAOJIK antibodies are further discussed below.

In preferred embodiments, a TAOJIK-interacting protein specifically binds a TAOJIK protein. In alternative preferred embodiments, a TAOJIK-modulating agent binds a TAOJIK substrate, binding partner, or cofactor.

Antibodies

In another embodiment, the protein modulator is a TAOJIK specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify TAOJIK modulators. The antibodies can also be used in dissecting the portions of the TAOJIK pathway responsible for various cellular responses and in the general processing and maturation of the TAOJIK.

Antibodies that specifically bind TAOJIK polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of TAOJIK polypeptide, and more preferably, to human TAOJIK. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

- Epitopes of TAOJIK which are particularly antigenic can be selected, for example, by routine screening of TAOJIK polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Natl. Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol. Immunol. 20:483-89;
- 5 Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence shown in any of SEQ ID NOs:16-18. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1} , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577).
- 10 Antibodies may be generated against crude cell extracts of TAOJIK or substantially purified fragments thereof. If TAOJIK fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of a TAOJIK protein. In a particular embodiment, TAOJIK-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject
- 15 polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

- The presence of TAOJIK-specific antibodies is assayed by an appropriate assay
- 20 such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding TAOJIK polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

- Chimeric antibodies specific to TAOJIK polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin
- 25 constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al., Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608;
- 30 Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized

antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

TAOJK-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells that express the targeted protein. (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in

association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml.

Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

Nucleic Acid Modulators

Other preferred TAOJIK-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit TAOJIK activity. Preferred nucleic acid modulators interfere with the function of the TAOJIK nucleic acid such as DNA replication, transcription, translocation of the TAOJIK RNA to the site of protein translation, translation of protein from the TAOJIK RNA, splicing of the TAOJIK RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the TAOJIK RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to a TAOJIK mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. TAOJIK-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiamidate

intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No.

5 5,235,033; and US Pat No. 5,378,841).

Alternative preferred TAOJIK nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to
10 the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998. Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952. (1999); Hammond, S. M., et al.,
15 Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene
20 expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous
25 clinical trials to be safe and effective (Milligan JF, et al, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL et al., Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, a TAOJIK-specific nucleic acid modulator is used in an assay to further elucidate the role of the TAOJIK in the beta-catenin pathway,
30 and/or its relationship to other members of the pathway. In another aspect of the invention, a TAOJIK-specific antisense oligomer is used as a therapeutic agent for treatment of beta-catenin-related disease states.

Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of TAOJK activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the TAOJK nucleic acid or protein. In general, secondary assays further assess the activity of a TAOJK modulating agent identified by a primary assay and may confirm that the modulating agent affects TAOJK in a manner relevant to the beta-catenin pathway. In some cases, TAOJK modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising a TAOJK polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. kinase activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates TAOJK activity, and hence the beta-catenin pathway. The TAOJK polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding),

transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of TAOJIK and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when TAOJIK-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the TAOJIK protein may be assayed by various known methods such as substrate processing (*e.g.* ability of the candidate TAOJIK-specific binding agents to function as negative effectors in TAOJIK-expressing cells), binding equilibrium constants (usually at least about 10^7 M⁻¹, preferably at least about 10^8 M⁻¹, more preferably at least about 10^9 M⁻¹), and immunogenicity (*e.g.* ability to elicit TAOJIK specific antibody in a heterologous host such as a mouse, rat, goat or rabbit). For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a TAOJIK polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The TAOJIK polypeptide can be full length or a fragment thereof that retains functional TAOJIK activity. The TAOJIK polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The TAOJIK polypeptide is preferably human TAOJIK, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of TAOJIK interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has TAOJIK-specific binding activity, and can be used to assess normal TAOJIK gene function.

Suitable assay formats that may be adapted to screen for TAOJIK modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput

and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, Curr Opin Chem Biol (1998) 2:597-603; Sundberg SA, Curr Opin Biotechnol 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved
5 fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.*, Selvin PR, Nat Struct Biol (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, Curr Opin Chem Biol (2000) 4:445-451).

10 A variety of suitable assay systems may be used to identify candidate TAOJIK and beta-catenin pathway modulators (*e.g.* U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

15 **Kinase assays.** In some preferred embodiments the screening assay detects the ability of the test agent to modulate the kinase activity of a TAOJIK polypeptide. In further embodiments, a cell-free kinase assay system is used to identify a candidate beta-catenin modulating agent, and a secondary, cell-based assay, such as an apoptosis or
20 hypoxic induction assay (described below), may be used to further characterize the candidate beta-catenin modulating agent. Many different assays for kinases have been reported in the literature and are well known to those skilled in the art (*e.g.* U.S. Pat. No. 6,165,992; Zhu et al., Nature Genetics (2000) 26:283-289; and WO0073469).

Radioassays, which monitor the transfer of a gamma phosphate are frequently used. For
25 instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from gamma -³³P ATP to a biotinylated peptide substrate; the substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, J Biomol Screen (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA
30 bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand.

Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor

in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133).

Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

Apoptosis assays. Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). An apoptosis assay system may comprise a cell that expresses a TAOJK, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether TAOJK function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express TAOJK relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the TAOJK plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et*

al., 1986, *Int. J. Cancer* 38, 369; Campana *et al.*, 1988, *J. Immunol. Meth.* 107, 79), or by other means.

Cell Proliferation may also be examined using [³H]-thymidine incorporation (Chen, J., 1996, *Oncogene* 13:1395-403; Jeoung, J., 1995, *J. Biol. Chem.* 270:18367-73).

- 5 This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [³H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter). Another proliferation assay uses the dye Alamar Blue (available from
- 10 Biosource International), which fluoresces when reduced in living cells and provides an indirect measurement of cell number (Voytik-Harbin SL *et al.*, 1998, *In Vitro Cell Dev Biol Anim* 34:239-46).

- Cell proliferation may also be assayed by colony formation in soft agar (Sambrook *et al.*, *Molecular Cloning*, Cold Spring Harbor (1989)). For example, cells transformed
- 15 with TAOJIK are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

- Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW *et al.* (1986) *Int J Radiat Biol Relat Stud Phys Chem Med* 49:237-55). Cells transfected with a TAOJIK may be stained with propidium iodide and evaluated in a flow
- 20 cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

- Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses a TAOJIK, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent
- 25 can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system such as a cell-free kinase assay system. A
- 30 cell proliferation assay may also be used to test whether TAOJIK function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express TAOJIK relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the TAOJIK plays a direct role in cell proliferation or cell cycle.

Angiogenesis. Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses a TAOJIK, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether TAOJIK function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express TAOJIK relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the TAOJIK plays a direct role in angiogenesis. U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434, among others.

Hypoxic induction. The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with TAOJIK in hypoxic conditions (such as with 0.1% O₂, 5% CO₂, and balance N₂, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses a TAOJIK, and that optionally has defective beta-catenin function (e.g. beta-catenin is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate beta-catenin modulating agents. In some embodiments

of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate beta-catenin modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether TAOJIK function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express TAOJIK relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the TAOJIK plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2x final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF, and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., Bioconjug Chem. 2001 May-Jun;12(3):346-53).

Tubulogenesis. Tubulogenesis assays monitor the ability of cultured cells, generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include Matrigel™ (Becton Dickinson), an extract of basement membrane proteins containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4° C and forms a solid gel at 37° C. Other suitable matrices comprise extracellular components such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, Nature Medicine 5:1418-1423). These assays have traditionally involved stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a candidate agent modulates. Moreover, we have found that different target genes respond differently to stimulation with different pro-angiogenic agents, including inflammatory angiogenic factors such as TNF-alpha. Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing a TAOJIK's response to a variety of factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF-alpha, ephrin, etc.

Cell Migration. An invasion/migration assay (also called a migration assay) tests the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, J Biol Chem 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for determining cell number. In another exemplary set up, cells are fluorescently labeled and

migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing a TAOJIK's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

Sprouting assay. A sprouting assay is a three-dimensional *in vitro* angiogenesis assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900 μ l of methocel-collagen solution and pipetted into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents are added after 30 min by pipetting 100 μ l of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to determine the cumulative sprout length per spheroid.

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the TAOJIK protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting TAOJIK-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also be used to test antibody modulators.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance TAOJIK gene expression, preferably mRNA expression. In general, expression analysis comprises comparing TAOJIK expression in like
5 populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express TAOJIK) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that
10 TAOJIK mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most
15 commonly detected with specific antibodies or antisera directed against either the TAOJIK protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve TAOJIK mRNA expression, may also be used to
20 test nucleic acid modulators.

Secondary Assays

Secondary assays may be used to further assess the activity of TAOJIK-modulating agent identified by any of the above methods to confirm that the modulating agent affects
25 TAOJIK in a manner relevant to the beta-catenin pathway. As used herein, TAOJIK-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with TAOJIK.

30 Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express TAOJIK) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate TAOJIK-modulating agent results in changes in the beta-catenin pathway in comparison to untreated (or mock- or placebo-

treated) cells or animals. Certain assays use "sensitized genetic backgrounds", which, as used herein, describe cells or animals engineered for altered expression of genes in the beta-catenin or interacting pathways.

5 ***Cell-based assays***

Cell based assays may detect endogenous beta-catenin pathway activity or may rely on recombinant expression of beta-catenin pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any
10 other efficacious means.

Animal Assays

A variety of non-human animal models of normal or defective beta-catenin pathway may be used to test candidate TAOJIK modulators. Models for defective beta-
15 catenin pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the beta-catenin pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, beta-catenin pathway activity is assessed by
20 monitoring neovascularization and angiogenesis. Animal models with defective and normal beta-catenin are used to test the candidate modulator's affect on TAOJIK in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin, collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid
25 Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the TAOJIK. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice
30 are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on TAOJIK is assessed via tumorigenicity assays. Tumor xenograft assays are known in the

art (see, e.g., Ogawa K et al., 2000, Oncogene 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the TAOJIK endogenously are injected in the flank, 1×10^5 to 1×10^7 cells per mouse in a volume of 100 μ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors maybe utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

In another preferred embodiment, tumorogenicity is monitored using a hollow fiber assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate modulator. Implanted cells are generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate period of time, generally around six days, the implanted samples are harvested for evaluation of the candidate modulator. Tumorogenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in soft agar, the capacity of the cells to recover and replicate *in vitro*, etc.

In another preferred embodiment, a tumorogenicity assay use a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as transgenic tumor assays. In a preferred application, tumor development in the transgenic model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell

carcinomas (Hanahan D, 1985, Nature 315:115-122; Parangi S et al, 1996, Proc Natl Acad Sci USA 93: 2002-2007; Bergers G et al, 1999, Science 284:808-812). An "angiogenic switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of small tumors (e.g., for a model of intervention), or during the growth of large and/or invasive tumors (e.g., for a model of regression). Tumorogenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

Diagnostic and therapeutic uses

Specific TAOJIK-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the beta-catenin pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the beta-catenin pathway in a cell, preferably a cell pre-determined to have defective or impaired beta-catenin function (e.g. due to overexpression, underexpression, or misexpression of beta-catenin, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates TAOJIK activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the beta-catenin function is restored. The phrase "function is restored", and equivalents, as used herein, means that the desired phenotype is achieved, or is brought closer to normal compared to untreated cells. For example, with restored beta-catenin function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal relative to untreated cells. The invention also provides methods for treating disorders or disease associated with impaired beta-catenin function by administering a therapeutically effective amount of a TAOJIK -modulating agent that modulates the beta-catenin pathway. The invention further provides methods for modulating TAOJIK function in a cell, preferably a cell pre-determined to have defective or impaired TAOJIK function, by administering a TAOJIK -modulating agent. Additionally, the invention provides a method for treating disorders or disease associated with impaired TAOJIK function by administering a therapeutically effective amount of a TAOJIK -modulating agent.

The discovery that TAOJIK is implicated in beta-catenin pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the beta-catenin pathway and for the identification of subjects having a predisposition to such diseases and disorders.

- 5 Various expression analysis methods can be used to diagnose whether TAOJIK expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47).
- 10 Tissues having a disease or disorder implicating defective beta-catenin signaling that express a TAOJIK, are identified as amenable to treatment with a TAOJIK modulating agent. In a preferred application, the beta-catenin defective tissue overexpresses a TAOJIK relative to normal tissue. For example, a Northern blot analysis of mRNA from
- 15 tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial TAOJIK cDNA sequences as probes, can determine whether particular tumors express or overexpress TAOJIK. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of TAOJIK expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).
- 20 Various other diagnostic methods may be performed, for example, utilizing reagents such as the TAOJIK oligonucleotides, and antibodies directed against a TAOJIK, as described above for: (1) the detection of the presence of TAOJIK gene mutations, or the detection of either over- or under-expression of TAOJIK mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of TAOJIK gene
- 25 product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by TAOJIK.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in TAOJIK expression, the method comprising: a) obtaining a biological sample from the patient; b) contacting

30 the sample with a probe for TAOJIK expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer, most preferably a cancer as shown in TABLE 2, or indicated as a result of immunohistochemistry analysis (Example VII). The probe may be either DNA or protein, including an antibody.

EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

5 I. C. elegans beta-catenin screen

The identification of mutants that suppress the cell adhesion defect of beta-catenin may lead to unique therapeutic targets that inhibit cell migration or metastasis. *hmp-2* was initially identified in an EMS screen for defects in body elongation during embryonic morphogenesis (see Costa *et al.*, (1998) The Journal of Cell Biology 1998, 141: 297-308).

- 10 The loss of function allele *hmp-2* (*zu364*) exhibits 99% embryonic lethality, with mutant embryos arresting during elongation and abnormal bulges forming on the dorsal side. About 1% of these embryos hatch to form viable lumpy larvae. The reduction of function allele *hmp-2* (*qm39*) yields viable larvae with a characteristic lumpy appearance. When grown at 15°C, approximately 92% (SD 3.9) of the L1 larvae show this lumpy phenotype,
- 15 with the penetrance of the phenotype decreasing as the animals molt and move through successive larval stages. For this screen, *hmp-2* (*qm39*) worms were soaked at 15°C in double stranded RNA (dsRNA) at the L4 larval stage and the progeny were scored as L1 larvae for modification of the adhesion defect. The screen protocol is described below.

- 20 1) *hmp-2* (*qm39*) animals were bleached and hatched on peptone free agarose plates to produce a synchronous population. Starved L1s were transferred to 10x peptone plates seeded with 750 µl OP50 (25% w/v in TB) and allowed to develop to the L4 larval stage.

- 25 2) dsRNA was dispensed in 6 µl aliquots into 96 well round bottom plates (Nunc #262162). L4 animals were collected by suspension in M9 buffer, washed 2x with M9 to remove any excess OP50, and dispensed in 2 µl aliquots into the RNA to a total worm density of 75-100 worms per well. As a control, multiple wells contained only RNA resuspension buffer (1x IM buffer).

- 3) Animals were soaked in dsRNA at 15°C for 24 hours.
- 30 4) Following dsRNA soaking, the animals were fed in the wells by addition of 25µl liquid NGM + 3% OP50. The animals were kept at 15°C and allowed to become gravid and lay progeny in the wells, which took approximately 72 hours. Food levels were monitored visually during maturation and more was added as needed.

5) Following maturation, animals from each well were plated onto individual 6cm peptone free agarose plates and placed at 15°C overnight.

6) Animals on each plate were scored visually under the dissecting microscope for modification of the lumpy phenotype. Scoring was performed qualitatively, with an increase in dead embryos scored as enhancement and an increase in wild type appearing animals scored as suppression of the defect.

7) Retests of interesting suppressor candidates followed the same protocol as the primary screen with certain modifications: several retests were performed for each suppressor, retested candidates were encoded so that they could be scored blindly, and retested candidates were scored quantitatively. Each plate was scored by counting 100 total objects. An object was defined as either an embryo or an L1 stage larva. Each object was scored as one of the following: a wildtype appearing animal, a lumpy appearing animal, or an unhatched embryo. Scores were represented as the percentage of wildtype appearing animals relative to all objects scored. Wildtype animals were defined as L1 larvae with smooth cuticles that did not have any sort of lumpy body morphology.

8) A confirmed suppressor was one that was ≥ 2 standard deviations away from the mean of the controls for at least 3 of the four retest experiments.

BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of the *C. elegans* modifiers. For example, representative sequences from TAOJIK, GI# 4759208 (SEQ ID NO:16), GI# 7705560 (SEQ ID NO:17) and GI#7243103 (SEQ ID NO:18) share 37%, 38%, and 38% amino acid identity, respectively, with the *C. elegans* T17E9.1.

Various domains, signals, and functional subunits in proteins were analyzed using the PSORT (Nakai K., and Horton P., Trends Biochem Sci, 1999, 24:34-6; Kenta Nakai, Protein sorting signals and prediction of subcellular localization, Adv. Protein Chem. 54, 277-344 (2000)), PFAM (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2), SMART (Ponting CP, et al., SMART: identification and annotation of domains from signaling and extracellular protein sequences. Nucleic Acids Res. 1999 Jan 1;27(1):229-32), TM-HMM (Erik L.L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology, p 175-182 Ed J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, CA: AAAI Press, 1998), and clust (Remm M, and Sonnhammer E. Classification of transmembrane protein families in the *Caenorhabditis elegans* genome and identification of human orthologs. Genome Res. 2000 Nov;10(11):1679-89) programs. For example, the kinase

(PFAM 00069) domain of TAOJIK from GI#s 4759208, 7705560, and 7243103 (SEQ ID NOs:16, 17, and 18, respectively) are located at approximately amino acid residues 28 to 281, 24 to 277, and 32 to 285, respectively.

Results:

- 5 Numbers shown in Table 1 are the percentage of wild type appearing animals relative to all animals (wild-type, lumpy and embryos) scored per experiment. Control replicates were performed for each retest (#1-4) and the standard deviation for each control is listed beneath the average.

10 Table 1

Gene ID	Retest #1	Retest #2	Retest #3	Retest #4	Mean	Deviation
T17E9.1 (kin-18)	29	23	36	10	24.5	11.0302614
Controls average	9.4	9.4	8.1	8.1		
Deviation	3.5	4.1	3.9	3.9		

II. RNAi of *C. elegans* T17E9.1

- 15 T17E9.1/kin-18 (the *C. elegans* ortholog of human kinases TAO1, JIK, and KIAA1361) was initially identified in an RNAi gene inactivation screen as suppressors of a beta-catenin (*hmp-2*) lumpy body mutant phenotype resulting from reduced beta-catenin function in cytoskeletal organization at cell-cell adhesive junctions (Example I). Subsequent testing demonstrated that RNAi inactivation of the *C. elegans* gene also partially suppressed an axin (*pry-1*) developmental arrest mutant phenotype that appears to result from increased beta-catenin (*bar-1*) and TCF (*pop-1*) transcriptional activation function.
- 20

III. High-Throughput In Vitro Fluorescence Polarization Assay

- 25 Fluorescently-labeled TAOJIK peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of TAOJIK activity.
- 30

IV. High-Throughput In Vitro Binding Assay.

³³P-labeled TAOJIK peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl₂, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate beta-catenin modulating agents.

V. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells containing the TAOJIK proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at $15,000 \times g$ for 15 min. The cell lysate is incubated with 25 μ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

VI. Kinase assay

A purified or partially purified TAOJIK is diluted in a suitable reaction buffer, e.g., 50 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10 μ g/ml). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by

increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100 μ l. The reaction is initiated by the addition of ^{33}P -gamma-ATP (0.5 $\mu\text{Ci/ml}$) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation (Mg^{2+} or Mn^{2+}) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

VII. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues were obtained from Impath, UC Davis, Clontech, Stratagene, Ardaïs, Genome Collaborative, and Ambion.

TAQMAN ANALYSIS. TaqMan analysis was used to assess expression levels of the disclosed genes in various samples.

RNA was extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/ μ l. Single stranded cDNA was then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan assay (Applied Biosystems, Foster City, CA) were prepared according to the TaqMan protocols, and the following criteria: a) primer pairs were designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis was performed using a 7900HT instrument.

Taqman reactions were carried out following manufacturer's protocols, in 25 μ l total volume for 96-well plates and 10 μ l total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis was prepared using a universal pool of human cDNA samples, which is a

mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data were normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples were compared with matched normal tissues from the same patient. A gene was considered overexpressed in a tumor when the level of expression of the gene was 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue was not available, a universal pool of cDNA samples was used instead. In these cases, a gene was considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type was greater than 2 times the standard deviation of all normal samples (i.e., $\text{Tumor} - \text{average}(\text{all normal samples}) > 2 \times \text{STDEV}(\text{all normal samples})$).

Results are shown in Table 2. Number of pairs of tumor samples and matched normal tissue from the same patient are shown for each tumor type. Percentage of the samples with at least two-fold overexpression for each tumor type is provided. A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

Table 2.

GI#	SE Q ID NO	# of Bre ast rs	# of Pai rs	# of Colo n rs	# of Pai rs	Head and Neck rs	# of Pai rs	# of Kidn ey rs	# of Pai rs	# of Lu ng rs	# of Pai rs	# of Ov ary rs	# of Pai rs	# of Uteru s rs	# of Pai rs	Prost ate rs	# of Pai rs	# of Skin rs	# of Pai rs
4759207	1	0%	19	12%	33	0%	8	21%	24	0%	21	8%	12	5%	19	8%	12	33%	3
7705559	5	5%	19	15%	33	13%	8	21%	24	5%	20	8%	12	5%	19	25%	12	0%	3
7243102	10	5%	19	12%	33	13%	8	4%	24	5%	20	9%	12	16%	19	17%	12	33%	3

IMMUNOHISTOCHEMICAL ANALYSIS. Immunohistochemistry was used to localize TAOJIK protein in human tissue sections according to known methods (Thomas Boenisch, ed. (2001) Handbook, Immunochemical Staining Methods, 3rd Edition, Dako Corporation, Carpinteria, CA). Antibody to TAOJIK GI#7243103 (SEQ ID NO:18) (rat TAO1, BD Biosciences; San Diego, CA) was used for immunohistochemistry against tissue arrays containing 20 normal and 19 tumor tissues. Tissue sections were pre-treated with heat antigen retrieval in citrate buffer and the antibody was used at 20 ug/ml. In normal tissues, expression was observed in smooth muscle, breast myoepithelial cells and kidney glomeruli. In tumor tissues, there was overexpression in stomach stromal sarcomas, breast, endometrial, and bladder cancers.

VIII. TAOJIK RNAi

RNA interference experiments were carried out to knock down expression of TAOJIKs using small interfering RNAs (siRNA, Elbashir et al, *supra*). These experiments were performed against TAOJIKs GI#s 4759207 (SEQ ID NO:1), 7705559 (SEQ ID NO:5), and 20559660 (SEQ ID NO:12). For each experiment, three different siRNAs (21mer, double stranded RNA oligos with 2 base 3' overhangs, obtained from Genset Oligos/Proligo France SAS, France; Dharmacon Research Inc., Lafayette, CO) were transfected into cell lines (available from American Type Culture Collection (ATCC), Manassas, VA) at 100 nM using oligofectamine (Invitrogen). A mock transfection reagent without siRNA was included with each experiment. Cells were incubated for 3 days at 37 degrees. At the end of each experiment, some wells of cells were harvested for protein extracts and used in western analysis and parallel cells were put through a BrdU ELISA to measure cell proliferation.

A. Effect of siRNA treatment of TAOJIKs in cells on beta-catenin. A549 lung cancer cells were grown in a 16 well slide (Nalge Nunc International; Rochester, NY). Cells were fixed with 4% paraformaldehyde and labeled with B-catenin antibody (Cell Signaling), and rhodamine-conjugated phalloidin (Cytoskeleton Inc.; Denver, CO) for actin staining.

Results:

B-catenin localization to the plasma membrane was decreased in cells transfected with siRNAi against all TAOJIKs. In addition, no changes in cell-cell interactions were observed.

B. Effect of siRNA treatment of TAOJIKs on cell proliferation in cancer cell lines. Cell lines A549 (lung cancer line), MDA-MB-231T (breast cancer line), SW480, SW620, HCT116 (colon cancer cell lines) were incubated for 3 days with 100nM duplex siRNA oligonucleotides, as described above. siRNA to luciferase and cyclin D1 were used as negative and positive controls, respectively. Cells in triplicate were incubated with 10% alamarBlue reagent (Biosource International; Camarillo, CA) according to standard protocol, to measure metabolism. Cells in triplicate were incubated with BrdU labeling reagent (Roche; Mannheim, Germany) according to standard protocol, to measure DNA synthesis in S phase cells.

10 **Results:**

A549 and MDA-MB-231T cells treated with siRNA against SEQ ID NOs:1, 5, and 12 showed approximately a 50% reduction in alamarBlue or BrdU values compared to mock transfection. Furthermore, SW480, SW620 and HCT116 cells treated with siRNA against SEQ ID NO:12 showed approximately 50% decrease in Alamar blue or BrdU values compared to mock transfection. Thus, siRNA treatment of TAOJIKs reduces cell proliferation in cancer cell lines.

C. Western blot analysis of effect of siRNA treatment of TAOJIKs on beta-catenin, actin, and E-cadherin expression in A549, SW620, and HCT116 cancer cell lines. Triplicate wells from the above experiments A and B were lysed in RIPA buffer (Boston BioProducts, Inc.; Ashland, MA) with protease inhibitors (Roche) and phosphatase inhibitors. Protein concentrations of cell lysates were determined by the BCA method (Pierce; Rockford, IL). Samples were run on a 4-12% gradient gel (Invitrogen; Carlsbad, CA), transferred to PVDF membrane (Bedford, MA), and probed with antibodies against SEQ ID NO:18 (rat TAO1, BD Biosciences; San Diego, CA); B-catenin (Cell Signaling Technology; Beverly, MA); actin (Accurate Chemical and Scientific Corporation; Westbury, NY); glyceraldehyde-3-phosphate dehydrogenase (Advanced Immuno Chemical, Inc.; Long Beach, CA); and E-cadherin (BD Biosciences).

25 **Results:**

In A549 cells, siRNA treatment of TAOJIKs SEQ ID NO:1 and SEQ ID NO:5 decreased beta-catenin, E-cadherin and actin expression in A549 cells. . KIAA1361 siRNA treatment of TAOJIK SEQ ID NO:12 decreased expression of SEQ ID NO:12 and E-cadherin.

Conclusion:

Taken together, the RNAi and western blot experiments, in addition to reducing cell proliferation, suggest a link between TAOJKs and beta-catenin function. Since the *C. elegans* T17E9.1 is also implicated in the beta catenin pathway, the link between

5 TAOJKs and the beta-catenin pathway provides compelling evidence of evolutionary functional conservation between these orthologs.

WHAT IS CLAIMED IS:

1. A method of identifying a candidate beta-catenin pathway modulating agent, said method comprising the steps of:
 - 5 (a) providing an assay system comprising a TAOJIK polypeptide or nucleic acid;
 - (b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
 - (c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as
 - 10 a candidate beta-catenin pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that express the TAOJIK polypeptide.
- 15 3. The method of Claim 2 wherein the cultured cells additionally have defective beta-catenin function.
4. The method of Claim 1 wherein the assay system includes a screening assay comprising a TAOJIK polypeptide, and the candidate test agent is a small molecule
- 20 modulator.
5. The method of Claim 4 wherein the assay is a kinase assay.
6. The method of Claim 1 wherein the assay system is selected from the group consisting
- 25 of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay comprising a TAOJIK polypeptide and the candidate test agent is an antibody.
- 30 8. The method of Claim 1 wherein the assay system includes an expression assay comprising a TAOJIK nucleic acid and the candidate test agent is a nucleic acid modulator.

9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.
10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.
- 5 11. The method of Claim 1 additionally comprising:
(d) administering the candidate beta-catenin pathway modulating agent identified in (c) to a model system comprising cells defective in beta-catenin function and, detecting a phenotypic change in the model system that indicates that the beta-catenin function is restored.
- 10 12. The method of Claim 11 wherein the model system is a mouse model with defective beta-catenin function.
13. A method for modulating a beta-catenin pathway of a cell comprising contacting a
15 cell defective in beta-catenin function with a candidate modulator that specifically binds to a TAOJIK polypeptide comprising an amino acid sequence selected from group consisting of SEQ ID NOs:16, 17, and 18 whereby beta-catenin function is restored.
14. The method of claim 13 wherein the candidate modulator is administered to a
20 vertebrate animal predetermined to have a disease or disorder resulting from a defect in beta-catenin function.
15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.
- 25 16. The method of Claim 1, comprising the additional steps of:
(d) providing a secondary assay system comprising cultured cells or a non-human animal expressing TAOJIK ,
(e) contacting the secondary assay system with the test agent of (b) or an agent
30 derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and
(f) detecting an agent-biased activity of the second assay system,

wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate beta-catenin pathway modulating agent,

5 and wherein the second assay detects an agent-biased change in the beta-catenin pathway.

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

10 18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

19. The method of Claim 18 wherein the non-human animal mis-expresses a beta-catenin pathway gene.

15 20. A method of modulating beta-catenin pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a TAOJIK polypeptide or nucleic acid.

20 21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the beta-catenin pathway.

22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

25 23. A method for diagnosing a disease in a patient comprising:
(a) obtaining a biological sample from the patient;
(b) contacting the sample with a probe for TAOJIK expression;
(c) comparing results from step (b) with a control;
30 (d) determining whether step (c) indicates a likelihood of disease.

24. The method of claim 23 wherein said disease is cancer.

25. The method according to claim 24, wherein said cancer is a cancer as shown in Table 2 as having >25% expression level.

26. The method of claim 24, wherein said cancer is stomach stromal sarcoma, breast,
5 endometrial, or bladder cancer.

SEQUENCE LISTING

<110> Exelixis, Inc.

<120> TAOJIKs AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE

<130> EX02-141

<150> 60/340,312

<151> 2001-12-13

<160> 18

<170> PatentIn version 3.1

<210> 1

<211> 4242

<212> DNA

<213> Homo sapiens

<400> 1

```

agaatttcaa atatcaggtt caggccccctg cgtgcaccag tatccgggggt tcattcccccg      60
ggcgttcaaa tatcggattc agtctccatc ccgttcagat attcgggggtt cagaccccac      120
aatcagaaat ccggaattcg gcagctgtcg ccctcgacga gggggaggac tggaccgcga      180
ggtcagatta ggttgtcacc cctcccccctc cagggggaggc ttcccggggcc cgcccctcag      240
gaagggcgaa agccgaggaa gaggtggcaa ggggaaaggt ctcttgccc ctctccctgc      300
ttggcagagc cgctggagga ccccaggcgg aagcggaggc gctggggcac catagtgacc      360
cctaccaggc caggccccac tctcaggggc ccagggggcc accatgccag ctggggggccg      420
ggccgggagc ctgaaggacc cagatgtggc tgagctcttc ttcaaggatg acccagaaaa      480
gctcttctct gacctccggg aaattggcca tggcagcttt ggagccgtat actttgcccg      540
ggatgtcccg aatagtgagg tgggtggccat caagaagatg tcctacagtg ggaagcagtc      600
caatgagaaa tggcaagaca tcatcaagga ggtgcgggtc ttacagaagc tccggcatcc      660
caacaccatt cagtaccggg gctgttacct gagggagcac acggcttggc tggtaatgga      720
gtattgcctg ggctcagctt ctgaccttct agaagtgcac aagaaacccc ttcaggaggt      780
agagatcgca gctgtgacct acggggcgct tcagggcctg gcatactctg actcccacaa      840
catgatccat agggatgtga aggctggaaa catcctgctg tcagagccag ggtagtgaa      900
gctaggggac tttggttctg cgtccatcat ggcacctgcc aactccttcg tgggcacccc      960
atactggatg gcacccgagg tgatcctggc catggatgag gggcagtacg atggcaaagt      1020
ggacgtctgg tccttgggga taacctgcat cgagctggct gaacggaaac caccgtctt      1080
taacatgaat gcgatgagt ccttatacca cattgcacag aacgaatccc ccgtgctcca      1140
gtcaggacac tggctctgagt acttccggaa tttgtcgac tcctgtcttc agaaaatccc      1200

```

tcaagacaga	ccaacctcag	aggttctcct	gaagcaccgc	tttgtgctcc	gggagcggcc	1260
acccacagtc	atcatggacc	tgatccagag	gaccaaggat	gccgtgcggg	agctggacaa	1320
cctgcagtac	cgcaagatga	agaagatcct	gttccaagag	gcacccaacg	gccctggtgc	1380
cgaggcccca	gaggaggaag	aggaggccga	gccctacatg	caccggggccg	ggactctgac	1440
cagcctcgag	agtagccact	cagtgccccag	catgtccatc	agcgcctcca	gccagagcag	1500
ctccgtcaac	agcctagcag	atgcctcaga	caacgaggaa	gaggaggagg	aggaggagga	1560
agaggaggag	gaggaagaag	gccctgaagc	ccgggagatg	gccatgatgc	aggaggggga	1620
gcacacagtc	acctctcaea	gtccattat	ccaccggctg	ccgggctctg	acaacctata	1680
tgatgacccc	taccagccag	agataacccc	cagccctctc	cagccgcctg	cagccccagc	1740
tcccacttcc	accacctctt	ctgcccgcgg	ccgggcctac	tgccgtaacc	gagaccactt	1800
tgccaccatc	cgaaccgcct	ccctggtcag	ccgtcagatc	caggagcatg	agcaggactc	1860
tgcgctgcgg	gagcagctga	gcggctataa	gcggatgcga	cgacagcacc	agaagcagct	1920
gctggccctg	gagtcacggc	tgaggggtga	acgggaggag	cacagtgcac	ggctgcagcg	1980
ggagcttgag	gcgcagcggg	ctggcttttg	ggcagaggca	gaaaagctgg	ccgggcggca	2040
ccaggccata	ggtgagaagg	aggcacgagc	tgcccaggcc	gaggagcggg	agttccagca	2100
gcacatcctt	gggcagcaga	agaaggagct	ggctgccctg	ctggaggcac	agaagcggac	2160
ctacaaactt	cgcaaggaac	agctgaagga	ggagctccag	gagaacccca	gcactcccaa	2220
gcgggagaag	gccgagtggc	tgctgcggca	gaaggagcag	ctccagcagt	gccaggcgga	2280
ggaggaagca	gggctgctgc	ggcggcagcg	ccagtacttt	gagctgcagt	gtcggcagta	2340
caagcgcaag	atgttgctgg	ctcggcacag	cctggaccag	gacctgctgc	gggaggacct	2400
gaacaagaag	cagaccagga	aggacttgga	gtgtgcaactg	ctgcttcggc	agcacgaggc	2460
cacgcgggag	ctggagctgc	ggcagctcca	ggcgtgcag	cgcacgcggg	ctgagctcac	2520
ccgcctgcag	caccagacgg	agctgggcaa	ccagctggag	tacaacaagc	ggcgtgagca	2580
agagttgcgg	cagaagcatg	cggcccaggt	tcgccagcag	cccaagagcc	tcaaactctaa	2640
ggagctgcag	atcaagaagc	agttccagga	gacgtgtaag	atccagactc	ggcagtacaa	2700
ggctctgcga	gcacacttgc	tggagaccac	gccccaaagct	cagcacaaga	gcctccttaa	2760
gcggctcaag	gaagagcaga	cccgcaagct	ggcgatcttg	gcggagcagt	atgaccagtc	2820
catctcagag	atgctcagct	cacaggcgct	gcggcttgat	gagaccaggg	aggcagagtt	2880
ccaggccctt	cggcagcagc	ttcaacagga	gctggagctg	ctcaacgctt	accagagcaa	2940
gatcaagatc	cgcacagaga	gccagcacga	gagggagctg	cgggagctgg	agcagagggt	3000
cgcgctgcgg	cgggcactgc	tggagcagcg	ggtggaagag	gagctgctgg	ccctgcagac	3060

```

aggacgctcc gagcgaatcc gcagtctgct tgagcggcag gcccgtagaga tcgaggcctt 3120
cgatgcggaa agcatgagggc tgggcttctc cagcatggct ctgggggggca tcccggctga 3180
agctgctgcc cagggctatc ctgctccacc cctgccccca gcctggccct cccgtcccgt 3240
tcccgtttct ggggcacact ggagccatgg cctcctcca ccaggcatgc cccctccagc 3300
ctggcgctcag ccgctctctgc tggctcccc aggcccccca aactggctgg ggccccccac 3360
acaaagtggg acaccccgtag gcggagccct gctgctgcta agaaacagcc cccagccctt 3420
gcggcgggca gcctcggggg gcagtggcag tgagaatgtg ggccccctg ctgccgcggt 3480
gcccgggccc ctgagccgca gcaccagtgt cgcttcccac atcctcaatg gttcttccca 3540
cttctattcc tgaggtgcag cggggaggag cagatgagct gggcagggca ggggtgggtg 3600
gagcctgacc ctggagggca ctgagctgga ggccctgca agggtagggg acaagatgta 3660
ggctccagct cccctcagac ctctcatct catgagcttc ttggggctgg ccagtggccc 3720
agggccagct tggcgataga tgctcaagg ctgcctggga gcccgcctc cctaccatgg 3780
tgccaggggt ctccctccgc cacctaggaa aggagggaga tgtgcgtgtc aaatattcat 3840
ctagtccctt gggggagggg aaggggtgggt ctagacatac tatattcaga gaactatact 3900
accctcacag tgaggccctc agacctgcca cagggcagag caggctctggg gcctgaggca 3960
gggagaatga gaggccacct tactggcagg aaggatcagg atggggcttt ggggtcagga 4020
tgctgggtc tcttccgta actgtctgac gtctgtgcc gtcttgtctt ttatcttttt 4080
tttttttttt taattgggat cagggtctgg gcggggaaac aagggaagga ccttgggaagg 4140
ggctgctccc aggcctgggg ggcagtcgtg ggagccctc tcagctgtgg ggctggcaca 4200
gagccccagg caagctttta ataaactgtt ggttattcta ac 4242

```

<210> 2

<211> 3069

<212> DNA

<213> Homo sapiens

<400> 2

```

gagaccggga cgagaccggg gctgtggtgc ggagagaggc tgagacggag aagaggagag 60
gcagagaggg cgcgggggacc gtcagcagca ccttagctac aatcgttcag ctattctcgg 120
aagagagaag ggagagggag gaggccgggg cgggagtggg ggctgtcacc ctccggacccc 180
ggcgtgagag gggccgtgcg gccggacgtc ctccgggtgg gccccagtc ggtggccgaa 240
gacctacagc tcaggccctt gggccccaaa ttccaggct ttgcccctcc tctttctca 300
gataccggg taacagtctt catagtccag atatccggga ctccgggtccc aacctctcta 360
aacctgggtc tctgtttcat agaatttcaa atatcaggtt caggccctg cgtgcaccag 420
tatccggggg tcattccccg ggcgttcaaa tatcggtatc agtctccatc ccgttcagat 480

```

attcgggggtt	cagaccccac	aatcagaaat	ccggaattcg	gcagctgtcg	ccctcgacga	540
gggggaggac	tggaccgcga	ggtcagatta	ggttgtcacc	ccctcccctc	caggggagggc	600
ttcccggggc	cgcctctcag	gaagggcgaa	agccgaggaa	gaggtggcaa	ggggaaaggt	660
ctccttgccc	ctctccctgc	ttggcagagc	cgctggagga	ccccaggcgg	aagcggagggc	720
gctggggcac	catagtgcac	cctaccaggc	caggcccccac	tctcagggcc	cccagggggcc	780
accatgccag	ctggggggcg	ggccgggagc	ctgaaggacc	cagatgtggc	tgagctcttc	840
ttcaaggatg	accagaaaa	gctcttctct	gacctccggg	aaattggcca	tggcagcttt	900
ggagccgtat	actttgcccg	ggatgtccgg	aatagtgcgg	tggtggccat	caagaagatg	960
tcctacagtg	ggaagcagtc	caatgagaaa	tggcaagaca	tcacaaagga	ggtgcgggttc	1020
ttacagaagc	tccggcatcc	caacaccatt	cagtaccggg	gctgttacct	gagggagcac	1080
acggcttggc	tggtaatgga	gtattgcctg	ggctcagctt	ctgaccttct	agaagtgcac	1140
aagaaacccc	ttcaggaggt	agagatcgca	gctgtgacct	acggggcgct	tcagggcctg	1200
gcatactctg	actcccacaa	catgatccat	agggatgtga	aggctggaaa	catcctgctg	1260
tcagagccag	ggttagtga	gctaggggac	tttggttctg	cgtccatcat	ggcacctgcc	1320
aactccttcg	tgggcacccc	atactggatg	gcacccgagg	tgatcctggc	catggatgag	1380
gggcagtacg	atggcaaagt	ggacgtctgg	tccttgggga	taacctgcat	cgagctggct	1440
gaacggaaac	caccgctctt	taacatgaat	gcgatgagtg	ccttatacca	cattgcacag	1500
aacgaatccc	ccgtgtcca	gtcaggacac	tggtctgagt	acttccggaa	ttttgtcgac	1560
tcctgtcttc	agaaaatccc	tcaagacaga	ccaacctcag	aggttctcct	gaagcaccgc	1620
tttgtgctcc	gggagcggcc	acccacagtc	atcatggacc	tgatccagag	gaccaaggat	1680
gccgtgcggg	agctggacaa	cctgcagtac	cgcaagatga	agaagatcct	gttccaagag	1740
gcaccaaacg	gccctgggtg	cgaggcccca	gaggaggaag	aggaggccga	gccctacatg	1800
caccggggcg	ggactctgac	cagcctcgag	agtagccact	cagtgccccg	catgtccatc	1860
agcgctccca	gccagagcag	ctccgtcaac	agcctagcag	atgcctcaga	caacgaggaa	1920
gaggaggagg	aggaggagga	agaggaggag	gaggaagaag	gccctgaagc	ccgggagatg	1980
gccatgatgc	aggaggggga	gcacacagtc	acctctcaca	gtccattat	ccaccggctg	2040
ccgggctctg	acaacctata	tgatgacccc	taccagccag	agataacccc	cagccctctc	2100
cagccgcctg	cagccccagc	tcccacttcc	accacctctt	ctgcccgcgg	ccgggcctac	2160
tgccgtaacc	gagaccactt	tgccaccatc	cgaaccgcct	ccctgggtcag	ccgtcagatc	2220
caggagcatg	agcaggactc	tgcgctgcgg	gagcagctga	gcggctataa	gcggatgcga	2280
cgacagcacc	agaagcagct	gctggccctg	gagtcacggc	tgaggggtga	acgggaggag	2340


```

cacagtgcac ggctgcagcg ggagcttgag ggcagcggg ctggctttgg ggcagaggca 2400
gaaaagctgg cccggcggca ccaggccata ggtgagaagg aggcacgagc tgcccaggcc 2460
gaggagcgga agttccagca gcacatcctt gggcagcaga agaaggagct ggctgccctg 2520
ctggaggcac agaagcggac ctacaaactt cgcaaggaa agctgaagga ggagctccag 2580
gagaacccca gcactcccaa gcgggagaag gccgagtggc tgctgcggca gaaggagcag 2640
ctccagcagt gccaggcgga ggaggaagca gggctgctgc ggcggcagcg ccagtacttt 2700
gagctgcagt gtcgccagta caagcgcaag atgttgctgg ctggcacag cctggaccag 2760
gacctgctgc gggaggacct gaacaagaag cagacccaga aggacttggg gtgtgcactg 2820
ctgcttcggc agcacgagcg cacgcgggag ctggagctgc ggcagctcca ggccgtgcag 2880
cgcacgcggg ctgagctcac ccgcctgcag caccagacgg agctgggcaa ccagctggag 2940
tacaacaagc ggcgtgagca agagtgcgg cagaagcatg cggcccagggt tcgccagcag 3000
cccaagagcc tcaaaaaaaaa gacaaacaca aataaaatat ctgagcggaa aaaaaaaaaa 3060
aaaaaaaaa 3069

```

<210> 3

<211> 3254

<212> DNA

<213> Homo sapiens

<400> 3

```

ctggaggacc ccaggcggaa gcggaggcgc tggggcacca tagtgacccc taccaggcca 60
ggccccactc tcaggggccc caggggccac catgccagct gggggccggg ccgggagcct 120
gaaggaccca gatgtggctg agctcttctt caaggatgac ccagaaaagc tcttctctga 180
cctccgggaa attggccatg gcagctttgg agccgtatac ttgcccggg atgtccggaa 240
tagtgagggtg gtggccatca agaagatgtc ctacagtggg aagcagtcca atgagaaatg 300
gcaagacatc atcaaggagg tgcggttctt acagaagctc cggcatccca acaccattca 360
gtaccggggc tgttacctga gggagcacac ggcttggtg gtaatggagt attgcctggg 420
ctcagcttct gaccttctag aagtgcacaa gaaacccctt caggaggtag agatcgagc 480
tgtgaccac ggggcgctt agggcctggc atatctgcac tcccacaaca tgatccatag 540
ggatgtgaag gctggaaaca tcctgctgtc agagccaggg ttagtgaagc taggggactt 600
tggttctgcg tccatcatgg cacctgccaa ctcttctgtg ggcaccccat actggatggc 660
accgaggtg atcctggcca tggatgaggg gcagtacgat ggcaaagtgg acgtctggtc 720
cttggggata acctgcacg agctggctga acggaaacca ccgtcttcta acatgaatgc 780
gatgagtgcc ttataccaca ttgcacagaa cgaatcccc gtgctccagt caggacactg 840

```

gtctgagtac	ttccggaatt	ttgtcgactc	ctgtcttcag	aaaatccctc	aagacagacc	900
aacctcagag	gttctcctga	agcaccgctt	tgtgctccgg	gagcggccac	ccacagtcac	960
catggacctg	atccagagga	ccaaggatgc	cgtgcgggag	ctggacaacc	tgcagtaccg	1020
caagatgaag	aagatcctgt	tccaagaggc	acccaacggc	cctggtgccg	aggccccaga	1080
ggaggaagag	gaggccgagc	cctacatgca	ccgggcccgg	actctgacca	gcctcgagag	1140
tagccactca	gtgcccagca	tgtccatcag	cgctccagc	cagagcagct	ccgtcaacag	1200
cctagcagat	gcctcagaca	acgaggaaga	ggaggaggag	gaggaggaag	aggaggagga	1260
ggaagaaggc	cctgaagccc	gggagatggc	catgatgcag	gagggggagc	acacagtcac	1320
ctctcacagc	tccattatcc	accggctgcc	gggctctgac	aacctatatg	atgaccacct	1380
ccagccagag	ataaccccca	gccctctcca	gccgcctgca	gccccagctc	ccacttccac	1440
cacctcttcc	gcccgcgcgc	gggcctactg	ccgtaaccga	gaccactttg	ccaccatccg	1500
aaccgcctcc	ctggtcagcc	gtcagatcca	ggagcatgag	caggactctg	cgctgcggga	1560
gcagctgagc	ggctataaag	ggatgcgacg	acagcaccag	aagcagctgc	tggccctgga	1620
gtcacggctg	aggggtgaac	gggaggagca	cagtgcacgg	ctgcagcggg	agcttgaggc	1680
gcagcgggct	ggctttgggg	cagaggcaga	aaagctggcc	cggcggcacc	aggccatagg	1740
tgagaaggag	gcacgagctg	cccaggccga	ggagcggaag	ttccagcagc	acatccttgg	1800
gcagcagaag	aaggagctgg	ctgccctgct	ggaggcacag	aagcggacct	acaaacttcg	1860
caaggaacag	ctgaaggagg	agctccagga	gaacccagc	actcccaagc	gggagaaggc	1920
cgagtggctg	ctgcggcaga	aggagcagct	ccagcagctc	caggcggagg	aggaagcagg	1980
gctgctgcgg	cggcagcgcc	agtactttga	gctgcagtgt	cgccagtaca	agcgcaagat	2040
gttgctggct	cggcacagcc	tggaccagga	cctgctgcgg	gaggacctga	acaagaagca	2100
gaccagaag	gacttggagt	gtgcaactgt	gcttcggcag	cacgaggcca	cgcggggagct	2160
ggagctgcgg	cagctccagg	ccgtgcagcg	cacgcgggct	gagctcacc	gcctgcagca	2220
ccagacggag	ctgggcaacc	agctggagta	caacaagcgg	cgtgagcaag	agttgcggca	2280
gaagcatgcg	gcccagggtc	gccagcagcc	caagagcctc	aaatctaagg	agctgcagat	2340
caagaagcag	ttccaggaga	cgtgtaagat	ccagactcgg	cagtacaagg	ctctgcgagc	2400
acacttgctg	gagaccacgc	ccaaagctca	gcacaagagc	ctccttaagc	ggctcaagga	2460
agagcagacc	cgcaagctgg	cgatcttggc	ggagcagtat	gaccagtcca	tctcagagat	2520
gctcagctca	caggcgctgc	ggcttgatga	gaccaggag	gcagagttcc	aggcccttcg	2580
gcagcagctt	caacaggagc	tggagctgct	caacgcttac	cagagcaaga	tcaagatccg	2640
cacagagagc	cagcacgaga	gggagctgcg	ggagctggag	cagagggtcg	cgctgcggcg	2700

ggcactgctg gagcagcggg tggaagagga gctgctggcc ctgcagacag gacgctccga 2760
 gcgaatccgc agtctgcttg agcggcaggc ccgtgagatc gaggccttcg atgcggaaag 2820
 catgaggctg ggcttctcca gcatggctct ggggggcatc ccggctgaag ctgctgcca 2880
 gggtatacct gctccacccc ctgccccagc ctggccctcc cgtcccgttc cccgttctgg 2940
 ggcacactgg agccatggcc ctctccacc aggcattgcc cctccagcct ggcgtcagcc 3000
 gtctctgctg gctccccag gcccccaaa ctggctgggg cccccacac aaagtgggac 3060
 acccctggc ggagccctgc tgctgctaag aaacagcccc cagcccctgc ggcgggcagc 3120
 ctgggggggc agtggcagtg agaattgtgg ccccctgct gccgcggtgc ccgggcccct 3180
 gagccgcagc accagtgtcg ctccacat cctcaatggt tcttcccact tctattcctg 3240
 aggtgcagcg ggga 3254

<210> 4
 <211> 4971
 <212> DNA
 <213> Homo sapiens

<400> 4
 aattcggcac gagctgagac ggagaagagg agaggcagag agggcgcggg gaccgtcagc 60
 agcaccttag ctacaatcgt tcagctatct tcggaagaga gaaggagag ggaggaggcc 120
 ggggcgggag tgggggctgt caccctcgga cccggcggtg agagggggcg tgccggcgga 180
 cgtcctcggg gtgggcccc agtcgggtggc cgaagacct cagctcaggc ccctgggtcc 240
 caaatttcca ggctttgccc ctctctcttt ctccagatac cgggtaacag tctcatagt 300
 ccagatatcc gggactcggg tcccaacctc tctaaacctg ggtctctgtt tcatagattt 360
 tcaaatatca ggttcaggcc cctgctgca ccagtatccg gggttcatc cccggggcgtt 420
 tcaaatatcg gattcagtt ccatcccggt cagatattcg gggttcagac cccacaatca 480
 gaaatccgga attcggcagc tgtcgccctc gacgaggggg aggactggac cgcgaggtca 540
 gattaggttg tcacccctc ccctccaggg gaggcttccc gggcccgccc ctccaggaagg 600
 gcgaaagccg aggaagagggt ggcaagggga aaggtctcct tgcccctctc cctgcttggc 660
 agagccgctg gaggacccca ggcggaagcg gaggcgctgg ggcaccatag tgaccctac 720
 caggccaggc cccactctca gggcccccag gggccaccat gccagctggg ggcggggccg 780
 ggagcctgaa ggaccagat gtggctgagc tcttcttcaa ggatgacca gaaaagctct 840
 tctctgacct ccgggaaatt ggccatggca gctttggagc cgtatacttt gccggggatg 900
 tccggaatag tgaggtggtg gccatcaaga agatgtccta cagtgggaag cagtccaatg 960
 agaaatggca agacatcatc aaggagggtgc ggttcttaca gaagctccgg catcccaaca 1020
 ccattcagta ccggggctgt tacctgaggg agcacacggc ttggctggta atggagtatt 1080

gcctgggctc agctttctgac cttctagaag tgcacaagaa accccttcag gaggtagaga 1140
 tcgcagctgt gacccacggg gcgcttcagg gcctggcata tctgcactcc cacaacatga 1200
 tccataggga tgtgaaggct ggaaacatcc tgctgtcaga gccaggggta gtgaagctag 1260
 gggacttttg ttctgcgtcc atcatggcac ctgccaaactc cttcgtgggc accccatact 1320
 ggatggcacc cgaggtgatc ctggccatgg atgaggggca gtacgatggc aaagtggacg 1380
 tctggtcctt ggggataacc tgcacgagc tggctgaacg gaaaccaccg ctctttaaca 1440
 tgaatgcgat gagtgcctta taccacattg cacagaacga atccccctg ctccagctag 1500
 gacactggtc tgagtacttc cggaattttg tcgactcctg tcttcagaaa atccctcaag 1560
 acagaccaac ctcagagggt ctcctgaagc accgctttgt gctccgggag cggccaccca 1620
 cagtcatcat ggacctgatc cagaggacca aggatgccgt gcgggagctg gacaacctgc 1680
 agtaccgcaa gatgaagaag atcctgttcc aagaggcacc caacggccct ggtgccgagg 1740
 ccccagagga ggaagaggag gccgagccct acatgcaccg ggccgggact ctgaccagcc 1800
 tcgagagtag ccactcagtg cccagcatgt ccatcagcgc ctccagccag agcagctccg 1860
 tcaacagcct agcagatgcc tcagacaacg aggaagagga ggaggaggag gaggaagagg 1920
 aggaggagga agaaggccct gaagcccggg agatggccat gatgcaggag ggggagcaca 1980
 cagtcacctc tcacagctcc attatccacc ggctgccggg ctctgacaac ctatatgatg 2040
 accctacca gccagagata acccccagcc ctctccagcc gcctgcagcc ccagctccca 2100
 cttccaccac ctcttcgcc cgccgccggg cctactgccg taaccgagac cactttgcca 2160
 ccatccgaac cgctccctg gtcagccgtc agatccagga gcatgagcag gactctgcgc 2220
 tgcgggagca gctgagcggc tataagcgga tgcgacgaca gcaccagaag cagctgctgg 2280
 ccctggagtc acggctgagg ggtgaacggg aggagcacag tgcacggctg cagcgggagc 2340
 ttgaggcgca gcggtctggc tttggggcag aggcagaaaa gctggcccgg cggcaccagg 2400
 ccataggtga gaaggaggca cgagctgccc aggccgagga gcggaagttc cagcagcaca 2460
 tccttgggca gcagaagaag gagctggctg ccctgctgga ggcacagaag cggacctaca 2520
 aacttcgcaa ggaacagctg aaggaggagc tccaggagaa cccagcact cccaagcggg 2580
 agaaggccga gtggctgctg cggcagaagg agcagctcca gcagtgccag gcggaggagg 2640
 aagcagggtc gctgcggcgg cagcgccagt actttgagct gcagtgtcgc cagtacaagc 2700
 gcaagatgtt gctggctcgg cacagcctgg accaggacct gctgcgggag gacctgaaca 2760
 agaagcagac ccagaaggac ttggagtgtg cactgctgct tcggcagcac gaggccacgc 2820
 gggagctgga gctgcggcag ctccaggccg tgcagcgcac gcgggctgag ctacccgcc 2880
 tgcagcacca gacggagctg ggcaaccagc tggagtacaa caagcggcgt gagcaagagt 2940

tgccgcagaa	gcatgcccc	caggttcgcc	agcagcccaa	gagcctcaa	gtacgtgcag	3000
gccagcgccc	cccgccctt	ccactccca	ttctggggc	tctgggcca	cccaacacag	3060
gcacccctat	agaacagcag	ccctgctcac	ctggccagga	ggcagtcctg	gaccaaagaa	3120
tgcttggcga	ggaggaggaa	gcagttggag	agagaaggat	tctgggaaag	gaaggggcca	3180
ctttggagcc	caagcagcag	aggattctgg	gggaagaatc	aggagccct	agtcccagtc	3240
cacaaaaaca	tgggagcctg	gttgatgagg	aagtttgggg	tctgcctgag	gagatagagg	3300
agcttagggg	gccctccctt	gtaccccagg	agaggagcat	tggtggccag	gaggaggctg	3360
ggacgtggag	cttggtgggg	aaggaggatg	agagtcttct	ggatgaggag	tttgagcttg	3420
gctgggtcca	gggcccagca	ctgactcccg	tccctgagga	ggaggaagaa	gaggaagagg	3480
gggctccgat	tgggaccct	agggatcctg	gagatgggtg	tccttcccc	gacatccctc	3540
ctgaaccccc	tccaacacac	ctgaggccct	gccctgccag	ccagctccct	ggactcctgt	3600
cccatggcct	cctggccggc	ctctcctttg	cagtgggggc	ctcctctggc	ctcctgcccc	3660
tcctgctgct	gctgctgctt	ccattgctgg	cagcccaggg	tgggggtggc	ctgcaggcag	3720
cgctgctggc	ccttgagggtg	gggctggtgg	gtctgggggc	ctcctacctg	ctcctttgta	3780
cagccctgca	cctgccctcc	agtcttttcc	tactcctggc	ccaggggtacc	gcactggggg	3840
ccgtcctggg	cctgagctgg	cgccgaggcc	tcattgggtgt	tccctgggc	cttgagctg	3900
cctggctctt	agcttgcca	ggcctagctc	tacctctggt	ggctatggca	gcggggggca	3960
gatgggtgcg	gcagcagggc	ccccgggtgc	gccggggcat	atctcgactc	tggttgcggg	4020
ttctgctgcg	cctgtcacc	atggccttcc	gggcccctgca	gggctgtggg	gctgtggggg	4080
accggggtct	gtttgcactg	taccccaaaa	ccaacaagga	tggcttccgc	agccgcctgc	4140
ccgtccctgg	gccccggcgg	cgtaatcccc	gcaccaccca	acaccatta	gctctgttgg	4200
caagggtctg	ggtcctgtgc	aagggtgga	actggcgtct	ggcacgggcc	agccagggtt	4260
tagcatccca	cttgcccccg	tgggccatcc	acacactggc	cagctggggc	ctgcttcggg	4320
gtgaacggcc	caccgaatc	ccccggtac	taccacgcag	ccagcgccag	ctagggcccc	4380
ctgcctccca	ccagccactg	ccagggactc	tagccggggc	gaggtcacgc	acccgccagt	4440
cccgggccct	gccccctgg	aggtagctga	ctccagccct	tccagcccaa	atctagagca	4500
ttgagcactt	tatctcccac	gactcagtga	agtttctcca	gtccctagtc	ctctcttttc	4560
accaccttc	ctcagtttgc	tcacttacc	caggcccagc	ccttcggacc	tctagacagg	4620
cagctcctc	agctgtggag	tccagcagtc	actctgtgtt	ctcctggcgc	tcctcccta	4680
agttattgct	gttcgcccgc	tgtgtgtgct	catcctcacc	ctcattgact	caggcctggg	4740
gccaggggtg	gtggagggtg	ggaagagtca	tgtttttttt	ctcctctttg	attttgtttt	4800

tctgtctccc ttccaacctg tccccttccc cccaccaaaa aaagaaaaag acaaacacaa 4860
 ataaaatatac tgagcggaac tgtgaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 4920
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 4971

<210> 5
 <211> 2958
 <212> DNA
 <213> Homo sapiens

<400> 5
 acggccatta ccaatcgcgga aaccaaagga ctgaagttat aaaagagaaa agagaagttt 60
 gctgctaaaa tgaatctgag caatatggaa tattttgtgc cacacacaaa aagggtactga 120
 agatttacc cccaaaaaaa attgtcaatg agaaataaag ctaactgata tcaaaaagca 180
 gagcctgctg tactggccat catgcgtaaa ggggtgctga aggaccaga gattgccgat 240
 ctattctaca aagatgatcc tgaggaaactt tttattgggt tgcatgaaat tggacatgga 300
 agttttggag cagtttattt tgctacaaat gctcacacca gtgaggtggt ggcaattaag 360
 aagatgtcct atagtgggaa gcagaccat gagaaatggc aagatattct taaggaagtt 420
 aaatttttac gacaattgaa gcatcctaact actattgagt acaaaggctg ttacttgaaa 480
 gaacacactg cttggttggt gatggaatat tgcttaggct cagcctctga tttattagaa 540
 gttcataaaa aaccacttca ggaagtggag atcgctgcca ttactcatgg agccttgcat 600
 ggactagcct acctacattc tcatgcattg attcataggg atattaaagc aggaaatatt 660
 cttctaacag agccaggta ggtaaaacta gctgattttg gatctgcttc aatggcttct 720
 cctgccaaact ccttctgtggg cacaccttac tggatggctc cagaagtgat cttagctatg 780
 gatgaaggac agtatgatgg gaaagttgat atttggcac ttggcatcac ttgtattgaa 840
 ttggcggaac ggaagccgcc ccttttcaac atgaatgcaa tgagtgcctt atatcacatt 900
 gcccagaatg actccccaac gttacagtct aatgaatgga cagactcctt taggagattt 960
 gttgattact gcttgcagaa aatacctcag gaaaggccaa catcagcaga actattaagg 1020
 catgactttg ttcgacgaga cgggccacta cgtgtcctca ttgacctcat acagaggaca 1080
 aaagatgcgg ttcgtgagct agataacct cagtaccgaa aaatgaaaaa aatacttttc 1140
 caagagacac ggaatggacc cttgaatgag tcacaggagg atgaggaaga cagtgaacat 1200
 ggaaccagcc tgaacaggga aatggacagc ctgggcagca accattccat tccaagcatg 1260
 tccgtaacat ggaaccagcc tgaacaggga aatggacagc ctgggcagca accattccat 1320
 tccaagcatg tccgtgtcat gatgcacgat gacgaaagca caatcaattc cagtcctcc 1380
 gtcgtgcata agaaagatca tgtattcata aggatgagg cgggccacgg cgatcccagg 1440

```

cctgagccgc ggccctaccca gtcagttcag agccaggccc tccactaccg gaacagagag 1500
cgcttttgcca cgatcaaatc agcatctttg gttacacgac agatccatga gcatgagcag 1560
gagaacgagt tgcgggaaca gatgtcaggt tataagcgga tgcggcgcca gcaccagaag 1620
cagctgatcg ccctggagaa caagctgaag gctgagatgg acgagcaccg cctcaagcta 1680
cagaaggagg tggagacgca tgccaacaac tcgtccatcg agctggagaa gctggccaag 1740
aagcaagtgg ctatcataga aaaggaggca aaggtagctg cagcagatga gaagaagttc 1800
cagcaacaga tcttggccca gcagaagaaa gatttgacaa ctttcttaga aagtcagaag 1860
aagcagtata agatttgtaa ggaaaaaata aaaggagaaa tgaatgagga ccatagcaca 1920
cccaagaaag agaagcaaga gcggatctcc aaacataaag agaacttgca gcacacacag 1980
gctgaagagg aagcccacct tctcactcaa cagagactgt actacgacaa aaattgtcgt 2040
ttcttcaagc ggaaaaataat gatcaagcgg cagcaggtgg agcagcagaa cattcgggag 2100
gaactaaata aaaagaggac ccagaaggag atggagcatg ccatgctaata ccggcagcag 2160
gagtccaccc gagagctaga gtacaggcag ctgcacacgt tacagaagct acgcatggat 2220
ctgatccgtt tacagcacca gacggaactg gaaaaccagc tggagtacaa taagaggcga 2280
gaaagagaac tgcacagaaa gcatgtcatg gaacttcggc aacagccaaa aaacttaaag 2340
gccatggaaa tgcaaatata aaaacagttt caggacactt gcaaagtaca gaccaaacag 2400
tataaagcac tcaagaatca ccagttggaa gttactccaa agaatgagca caaaacaatc 2460
ttaaagacac tgaaagatga gcagacaaga aaacttgcca ttttggcaga gcagtatgaa 2520
cagagtataa atgaaatgat ggcctctcaa gcgttacggc tagatgaggc tcaagaagca 2580
gaatgccagg ccttgaggct acagctccag caggaaatgg agctgctcaa cgctaccag 2640
agcaaaatca agatgcaaac agaggcacia catgaacgtg agctccagaa gctagagcag 2700
agagtgtctc tgcgcagagc acaccttgag cagaagattg aagaggagct ggctgccctt 2760
cagaaggaac gcagcgagag aataaagaac ctattggaaa ggcaagagcg agagattgaa 2820
acttttgaca tggagagcct cagaatggga tttgggaatt tggttacatt agattttcct 2880
aaggaggact acagatgaga ttaaattttt ttccatttac aaaaaaaaaa aaaaaaaaaa 2940
aaaaaaaaaa aaaaaaaa 2958

```

<210> 6

<211> 2897

<212> DNA

<213> Homo sapiens

<400> 6

```

caaaggactg aagttataaa agagaaaaga gaagtttgct gctaaatga atctgagcaa 60
tatggaatat tttgtgccac acacaaaaag gtactgaaga ttaccccc aaataaatt 120

```

gtcaatgaga aataaagcta actgatatca aaaagcagag cctgctctac tggccatcat	180
gcgtaaaggg gtgctgaagg acccagagat tgccgatcta ttctacaaag atgatacctga	240
ggaacttttt attggtttgc atgaaattgg acatggaagt tttggagcag tttatttttgc	300
tacaaatgct cacaccagtg aggtggtggc aattaagaag atgtcctata gtgggaagca	360
gacccatgag aaatggcaag atattcttaa ggaagttaaa tttttacgac aattgaagca	420
tcctaatact attgagtaca aaggctgtta cttgaaagaa cacactgctt ggttggtgat	480
ggaatattgc ttaggctcag cctctgattt attagaagtt cataaaaaac cacttcagga	540
agtggagatc gctgccatta ctcatggagc cttgcatgga ctgcctacc tacattctca	600
tgcattgatt catagggata ttaaagcagg aaatattctt ctaacagagc caggtcaggt	660
aaaactagct gattttggat ctgcttcaat ggcttctcct gccaaactcct tcgtgggcac	720
accttactgg atggctccag aggtgatctt agctatggat gaaggacagt atgatgggaa	780
agttgatatt tggtcacttg gcatcacttg tattgaattg gcggaacgga agccgcccct	840
tttcaacatg aatgcaatga gtgccttata tcacattgcc cagaatgact cccaacggt	900
acagtctaataaat gaatggacag actccttttag gagatttggt gattactgct tgcagaaaat	960
acctcaggaa aggccaacat cagcagaact attaaggcat gactttgttc gacgagaccg	1020
gccactacgt gtcctcattg acctcataca gaggacaaaa gatgcagttc gtgagctaga	1080
taacctacag taccgaaaaa tgaaaaaaat acttttccaa gagacacgga atggaccctt	1140
gaatgagtca caggaggatg aggaagacag tgaacatgga accagcctga acagggaaat	1200
ggacagcctg ggcagcaacc attccattcc aagcatgtcc gtgagcacag gcagccagag	1260
cagcagtgtg aacagcatgc aggaagtcac ggacgagagc agttccgaac ttgtcatgat	1320
gcacgatgac gaaagcacaa tcaattccag ctccctccgtc gtgcataaga aagatcatgt	1380
attcataagg gatgaggcgg gccacggcga tcccaggcct gagccgcggc ctaccagtc	1440
agttcagagc caggccctcc actaccggaa cagagagcgc tttgccacga tcaaatcagc	1500
atctttgggtt acacgacaga tccatgagca tgagcaggag aacgagttgc gggaacagat	1560
gtcaggttat aagcggatgc ggcgccagca ccagaagcag ctgatcgccc tggagaacaa	1620
gctgaaggct gagatggacg agcaccgcct caagctacag aaggaggtgg agacgcacgc	1680
caacaactcg tccatcgagc tggagaagct ggccaagaag caagtggcta tcatagaaaa	1740
ggaggcacaag gtagctgcag cagatgagaa gaagttccag caacagatct tggcccagca	1800
gaagaaagat ttgacaactt tcttagaaaag tcagaagaag cagtataaga tttgtaagga	1860
aaaaataaaa gaggaaatga atgaggacca tagcacaccc aagaaagaga agcaagagcg	1920
gatctccaaa cataaagaga acttgcagca cacacaggct gaagaggaag cccaccttct	1980

cactcaacag agactgtact acgacaaaaa ttgtcgtttc ttcaagcgga aaataatgat 2040
 caagcggcac gaggtggagc agcagaacat tcggggaggaa ctaaataaaa agaggaccca 2100
 gaaggagatg gagcatgccā tgctaataccg gcacgacgag tccacccgag agctagagta 2160
 caggcagctg cacacgttac agaagctacg catggatctg atccgtttac agcaccagac 2220
 ggaactggaa aaccagctgg agtacaataa gaggcgagaa agagaactgc acagaaagca 2280
 tgtcatggaa cttcggcaac agccaaaaaa cttaaaggcc atggaaatgc aaattaaaaa 2340
 acagtttcag gacacttgca aagtacagac caaacagtat aaagcactca agaatcacca 2400
 gttggaagtt actccaaaga atgagcacaā aacaatctta aagacactga aagatgagca 2460
 gacaagaaaa cttgccattt tggcagagca gtatgaacag agtataaatg aaatgatggc 2520
 ctctcaagcg ttacggctag atgaggctca agaagcagaa tgccaggcct tgaggctaca 2580
 gctccagcag gaaatggagc tgctcaacgc ctaccagagc aaaatcaaga tgcaaacaga 2640
 ggcacaacat gaacgtgagc tccagaagct agagcagaga gtgtctctgc gcagagcaca 2700
 ccttgagcag aagattgaag aggagctggc tgcccttcag aaggaacgca gcgagagaat 2760
 aaagaaccta ttggaaaggc aagagcgaga gattgaaact tttgacatgg agagcctcag 2820
 aatgggattt gggaatttgg ttacattaga ttttcttaag gaggactaca gatgagatta 2880
 aattttttgc catttac 2897

<210> 7

<211> 3148

<212> DNA

<213> Homo sapiens

<400> 7

ggcacgaggg tggcgccggg cggcgggggtc ctgcgtggag agtgggacgc aacgccgaga 60
 ccgcgagcag aggctgcgca cagccggatc cggcactcag cgaccggacc caaggatccg 120
 ccggggaaca agccacagga gagcgactca ggaacaagtg tgggagagga agcggcgggcg 180
 gcggcgccgg gcccggggggt ggtgacagca ggtctgaggt tgcatacataa atacaaagga 240
 ctgaagttat aaaagagaaa agagaagttt gctgctaaaa tgaatctgag caatatggaa 300
 tattttgtgc cacacacaaa aaggtactga agattttacc cccaaaaaaa attgtcaatg 360
 agaaataaag ctaactgata tcaaaaagca gagcctgctc tactggccat catgcgtaaa 420
 ggggtgctga aggaccagga gattgccgat ctattctaca aagatgatcc tgaggaactt 480
 tttattgggt tgcatgaaat tggacatgga agttttggag cagttttatt tgctacaaat 540
 gctcacacca gtgaggtggg ggcaattaag aagatgtcct atagtgggaa gcagacccat 600
 gagaaatggc aagatattct taaggaagtt aaatttttac gacaattgaa gcataccta 660

actattgagt acaaaggctg ttacttgaaa gaacacactg cttgggttggg gatggaatat	720
tgcttaggct cagcctctga tttattagaa gttcataaaa aaccacttca ggaagtggag	780
atcgctgcca ttactcatgg agccttgcat ggactagcct acctacattc tcatgcattg	840
attcataggg atattaaagc aggaaatatt cttctaacag agccagggtca ggtaaaacta	900
gctgattttg gatctgcttc aatggcttct cctgccaaact ccttcgtggg cacaccttac	960
tggatggctc cagagggtgat cttagctatg gatgaaggac agtatgatgg gaaagttgat	1020
atttggtcac ttggcatcac ttgtattgaa ttggcggaac ggaagccgcc ccttttcaac	1080
atgaatgcaa tgagtgcctt atatcacatt gcccagaatg actccccaac gttacagtct	1140
aatgaatgga cagactcctt taggagattt gttgattact gcttgacaga aatacctcag	1200
gaaaggccaa catcagcaga actattaagg catgactttg ttcgacgaga ccggccacta	1260
cgtgtcctca ttgacctcat acagaggaca aaagatgcag ttcgtgagct agataaccta	1320
cagtaccgaa aaatgaaaaa aatacttttc caagagacac ggaatggacc cttgaatgag	1380
tcacaggagg atgaggaaga cagtgaacat ggaaccagcc tgaacaggga aatggacagc	1440
ctgggcagca accattccat tccaagcatg tccgtgagca caggcagcca gagcagcagt	1500
gtgaacagca tgcaggaagt catggacgag agcagttccg aacttgatcat gatgcacgat	1560
gacgaaagca caatcaattc cagctcctcc gtcgtgcata agaaagatca tgtattcata	1620
agggatgagg cggggccacgg cgatcccagg cctgagccgc ggcctacca gtcagttcag	1680
agccaggccc tccactaccg gaacagagag cgctttgcca cgatcaaata agcatctttg	1740
gttacacgac agatccatga gcatgagcag gagaacgagt tgcgggaaca gatgtcaggt	1800
tataagcgga tgcggcgcca gcaccagaag cagctgatcg ccctggagaa caagctgaag	1860
gctgagatgg acgagcaccg cctcaagcta cagaaggagg tggagacgca tgccaacaac	1920
tcgtccatcg agctggagaa gctggccaag aagcaagtgg ctatcataga aaaggaggca	1980
aaggtagctg cagcagatga gaagaagttc cagcaacaga tcttggccca gcagaagaaa	2040
gatttgacaa ctttcttaga aagtcagaag aagcagtata agatttgtaa ggaaaaata	2100
aaaggagaaa tgaatgagga ccatagcaca cccaagaaag agaagcaaga gcggatctcc	2160
aaacataaag agaacttgca gcacacacag gctgaagagg aagcccacct tctcactcaa	2220
cagagactgt actacgacaa aaattgtcgt ttcttcaagc ggaaaataat gatcaagcgg	2280
cacgaggtgg agcagcagaa cattcggggag gaactaaata aaaaggaggac ccagaaggag	2340
atggagcatg ccatgctaata ccggcacgac gagtccaccc gagagctaga gtacaggcag	2400
ctgcacacgt tacagaagct acgcatggat ctgatccgtt tacagcacca gacggaactg	2460
gaaaaccagc tggagtacaa taagaggcga gaaagagaac tgcacagaaa gcatgtcatg	2520

gaacttcggc aacagccaaa aaacttaaag gccatggaaa tgcaaattaa aaaacagttt 2580
 caggacactt gcaaagtaca gaccaaacag tataaagcac tcaagaatca ccagttggaa 2640
 gttactccaa agaatgagca caaaacaatc ttaaagacac tgaaagatga gcagacaaga 2700
 aaacttgcca ttttggcaga gcagtatgaa cagagtataa atgaaatgat ggcctctcaa 2760
 gcgttacggc tagatgaggc tcaagaagca gaatgccagg ccttgaggct acagctccag 2820
 caggaaatgg agctgctcaa cgcctaccag agcaaaatca agatgcaaac agaggcaciaa 2880
 catgaacgtg agctccagaa gctagagcag agagtgtctc tgcgcagagc acaccttgag 2940
 cagaagattg aagaggagct ggctgccctt cagaaggaac gcagcgagag aataaagaac 3000
 ctattggaaa ggcaagagcg agagattgaa acttttgaca tggagagcct cagaatggga 3060
 tttgggaatt tggttacatt agattttcct aaggaggact acagatgaga ttaaattttt 3120
 tgccatttac aaaaaaaaaa aaaaaaaaaa 3148

<210> 8

<211> 4188

<212> DNA

<213> Homo sapiens

<400> 8

gccggggaac aagccacagg agagcgactc aggaacaagt gtgggagagg aagcggcggc 60
 ggcggcgccg ggcccggggg tggtagacagc aggtctgagg ttgcatcata aatacaaagg 120
 actgaagtta taaaagagaa aagagaagtt tgctgctaaa atgaatctga gcaatatgga 180
 atattttgtg ccacacacaa aaagggtactg aagatttacc ccccaaaaaa aattgtcaat 240
 gagaaataaa gctaactgat atcaaaaagc agagcctgct ctactggcca tcatgcgtaa 300
 aggggtgctg aaggaccag agattgccga tctattctac aaagatgatc ctgaggaact 360
 ttttattggg ttgcatgaaa ttggacatgg aagttttgga gcagtattatt ttgetacaaa 420
 tgctcacacc aatgaggtgg tggcaattaa gaagatgtcc tatagtggga agcagaccca 480
 tgagaaatgg caagatatct ttaaggaagt taaattttta cgacaattga agcatcctaa 540
 tactattgag taaaaaggct gttacttgaa agaacacact gcttggttgg tgatggaata 600
 ttgcttaggc tcagcctctg atttattaga agttcataaa aaaccacttc aggaagtgga 660
 gatcgctgcc attactcatg gagccttgca tggactagcc tacctacatt ctcatgcatt 720
 gattcatagg gatattaaag caggaaatat tcttctaaca gagccaggtc aggtaaaact 780
 agctgatttt ggatctgctt caatggcttc tcttgccaac tccttcgtgg gcacacctta 840
 ctggatggct ccagaggtga tcttagctat ggatgaagga cagtatgatg ggaaagtga 900
 tatttggtca cttggcatca cttgtattga attggcggaa cggaagccgc cccttttcaa 960
 catgaatgca atgagtgcct tatatcacat tgcccagaat gactccccaa cgttacagtc 1020

taatgaatgg acagactcct ttaggagatt tgttgattac tgcttgacaga aaataacctca	1080
ggaaaggcca acatcagcag aactattaag gcatgacttt gttcgacgag accggccact	1140
acgtgtcctc attgacctca tacagaggac aaaagatgca gttcgtgagc tagataacct	1200
acagtaccga aaaatgaaaa aaataactttt ccaagagaca cggaatggac ccttgaatga	1260
gtcacaggag gatgaggaag acagtgaaca tggaaccagc ctgaacaggg aaatggacag	1320
cctgggcagc aaccattcca ttccaagcat gtccgtgagc acaggcagcc agagcagcag	1380
tgtgaacagc atgcaggaag tcatggacga gagcagttcc gaacttgtca tgatgcacga	1440
tgacgaaagc acaatcaatt ccagctcctc cgtcgtgcat aagaaagatc atgtattcat	1500
aagggatgag gcgggccacg gcgatcccag gcctgagccg cggcctaccc agtcagttca	1560
gagccaggcc ctccactacc ggaacagaga gcgctttgcc acgatcaa atcagcatcttt	1620
ggttacacga cagatccatg agcatgagca ggagaacgag ttgcgggaac agatgtcagg	1680
ttataagcgg atgcggcgcc agcaccagaa gcagctgac gccctggaga acaagctgaa	1740
ggctgagatg gacgagcacc gcctcaagct acagaaggag gtggagacgc atgccacaa	1800
ctcgtccatc gagctggaga agctggccaa gaagcaagtg gctatcatag aaaaggaggc	1860
aaaggtagct gcagcagatg agaagaagtt ccagcaacag atcttggccc agcagaagaa	1920
agatttgaca actttcttag aaagtcagaa gaagcagtat aagatttgta aggaaaaaat	1980
aaaagaggaa atgaatgagg accatagcac acccaagaaa gagaagcaag agcggatctc	2040
caaacataaa gagaacttgc agcacacaca ggctgaagag gaagcccacc ttctcactca	2100
acagagactg tactacgaca aaaattgtcg tttcttcaag cggaaaataa tgatcaagcg	2160
gcacgaggtg gagcagcaga acattcggga ggaactaaat aaaaagagga cccagaagga	2220
gatggagcat gccatgctaa tccggcacga cgagtccacc cgagagctag agtacaggca	2280
gctgcacacg ttacagaagc tacgcatgga totgatccgt ttacagcacc agacggaact	2340
ggaaaaccag ctggagtaca ataagaggcg agaaagagaa ctgcacagaa agcatgtcat	2400
ggaacttcgg caacagccaa aaaacttaaa ggccatggaa atgcaaatta aaaaacagtt	2460
tcaggacact tgcaaagtac agaccaaaca gtataaagca ctcaagaatc accagttgga	2520
agttactcca aagaatgagc acaaaacaat cttaaagaca ctgaaagatg agcagacaag	2580
aaaacttgcc attttggcag agcagtatga acagagtata aatgaaatga tggcctctca	2640
agcgttacgg ctagatgagg ctcaagaagc agaatgccag gccttgaggc tacagctcca	2700
gcaggaaatg gagctgctca acgcctacca gagcaaaatc aagatgcaaa cagaggcaca	2760
acatgaacgt gagctccaga agctagagca gagagtgtct ctgcgcagag cacaccttga	2820
gcagaagatt gaagaggagc tggctgcct tcagaaggaa cgagcgaga gaataaagaa	2880

cctattggaa aggcaagagc gagagattga aacttttgac atggagagcc tcagaatggg 2940
 atttgggaat ttggttacat tagattttcc taaggaggac tacagatgag attaaatttt 3000
 ttgccattta caaaaaaaaa aaaaaaaga aaacagaaaa aaattcagac cctgcaaaac 3060
 cacattcccc attttaacgg gcgttgctct cactctctct ctctcttact cttactgaca 3120
 tcgtgtcgga ctagtgcctg tttattctta ctccatcagg ggcccccttc ctccccccgt 3180
 gtcaactttc agtgcctggc aaaacctggc cgtctcttct attcacagta cacgtcacag 3240
 tattgatgtg attcaaaatg tttcagtga aactttggag acagttttaa caaaaccaat 3300
 aaaccaacaa caaaaaaagt ggatgtatat tgctttaagc aatcactcat taccaccaat 3360
 ctgtgaaagt aaagcaaaaa ataataataa taaatgccaa gggggagaga gacacaatat 3420
 ccgcagcctt acaccttaac tagctgctgc attattttat tttattttat ttttttggtta 3480
 tttattcatc aggaataaaa aaaacaaagt tttattaaag attgaaaatt tgatacattt 3540
 tacagaaact aattgtgatg tacatatcag tggtgacata ttattacttt tttggggacg 3600
 ggggtgggtg ggggtgaagag atcttgtgat tttagactgc tgcagagtta acttgtctca 3660
 gcatactctga tgtatcataa tcatttctgc tgtgcagagg agggatacac ttaggggctc 3720
 acagatccca gtagcacaat tgggcttttg caaatgggta ttttgtgtat agaggaattt 3780
 aaggagaggt attacttatt ttcattattg attttaactg tttctcgat caaatttttt 3840
 aacttcttct tcgtgttctt cccacctcc ttccttttcc agttcagtat ttggagttca 3900
 acactgtctc tcaatcagat catctggatc tttttcttta tctcccttcc ccttccctaag 3960
 tcccatttct tggtcataaa tattgcatta ttcacacttt caaactgtgt attttcttac 4020
 aataaaaaat gatgaaaaaa aaaaaggctt tacttctttt gcatgcactt taaaaacaaa 4080
 acaaaacatt tttcaggttc caaggaagag catgataact gtcagagctt ttaattatat 4140
 ttgtaaataa aagtgttcat cacaaaaaaa aaaaaaaaaa aaaaaaaa 4188

<210> 9

<211> 2807

<212> DNA

<213> Homo sapiens

<400> 9

gtactgaaga tttaccccc aaaaaaatt gtcaatgaga aataaagcta actgatatca 60
 aaaagcagag cctgctctac tggccatcat gcgtaaaggg gtgctgaagg acccagagat 120
 tgccgatcta ttctacaaag atgatcctga ggaacttttt attggtttgc atgaaattgg 180
 acatggaagt tttggagcag tttattttgc tacaaatgct cacaccagtg aggtgggtggc 240
 aattaagaag atgtcctata gtgggaagca gacccatgag aaatggcaag atattcttaa 300

ggaagttaaa tttttacgac aattgaagca tcctaatact attgagtaca aaggctgtta 360
 cttgaaagaa cacactgctt ggttggtgat ggaatattgc ttaggctcag cctctgattt 420
 attagaagtt cataaaaaac cacttcagga agtggagatc gctgccatta ctcatggagc 480
 cttgcatgga ctagectacc tacattctca tgcattgatt catagggata ttaaagcagg 540
 aaatattctt ctaacagagc caggtcaggt aaaactagct gattttggat ctgcttcaat 600
 ggcttctcct gccaaactcct tcgtgggcac accttactgg atggctccag aggtgatctt 660
 agctatggat gaaggacagt atgatgggaa agttgatatt tggtcacttg gcatcacttg 720
 tattgaattg gcggaacgga agccgccctt tttcaacatg aatgcaatga gtgccttata 780
 tcacattgcc cagaatgact ccccaacgtt acagtctaataaat gaatggacag actcctttag 840
 gagatttgtt gattactgct tgcagaaaat acctcaggaa aggccaacat cagcagaact 900
 attaaggcat gactttgttc gacgagaccg gccactacgt gtcctcattg acctcataca 960
 gaggacaaaa gatgcagttc gtgagctaga taacctacag taccgaaaaa tgaaaaaaat 1020
 acttttccaa gagacacgga atggaccctt gaatgagtca caggaggatg aggaagacag 1080
 tgaacatgga accagcctga acagggaaat ggacagcctg ggcagcaacc attccattcc 1140
 aagcatgtcc gtgagcacag gcagccagag cagcagtgtg aacagcatgc aggaagtcatt 1200
 ggacgagagc agttccgaac ttgtcatgat gcacgatgac gaaagcaciaa tcaattccag 1260
 ctctccgctc gtgcataaga aagatcatgt attcataagg gatgaggcgg gccacggcga 1320
 tcccaggcct gagccgcggc ctaccagtc agttcagagc caggccctcc actaccggaa 1380
 cagagagcgc tttgccacga tcaaatacagc atctttggtt acacgacaga tccatgagca 1440
 tgagcaggag aacgagttgc gggaacagat gtcaggttat aagcggatgc ggcgccagca 1500
 ccagaagcag ctgatcgccc tggagaacaa gctgaaggct gagatggacg agcaccgcct 1560
 caagctacag aaggaggtgg agacgcatgc caacaactcg tccatcgagc tggagaagct 1620
 ggccaagaag caagtggcta tcatagaaaa ggaggcaaag gtagctgcag cagatgagaa 1680
 gaagtccag caacagatct tggcccagca gaagaaagat ttgacaactt tcttagaaaag 1740
 tcagaagaag cagtataaga tttgtaagga aaaaataaaa gaggaatga atgaggacca 1800
 tagcacacc aagaaagaga agcaagagcg gatctccaaa cataaagaga acttgcagca 1860
 cacacaggct gaagaggaag cccaccttct cactcaacag agactgtact acgacaaaaa 1920
 ttgtcgtttc ttcaagcgga aaataatgat caagcggcac gaggtggagc agcagaacat 1980
 tggggaggaa ctaaataaaa agaggaccca gaaggagatg gagcatgcca tgctaattcg 2040
 gcacgacgag tccacccgag agctagagta caggcagctg cacacgttac agaagctacg 2100
 catggatctg atccgtttac agcaccagac ggaactggaa aaccagctgg agtacaataa 2160

gaggcgagaa agagaactgc acagaaagca tgtcatggaa cttcggcaac agccaaaaaa 2220
 cttaaaggcc atggaaatgc aaattaaaaa acagtttcag gacacttgca aagtacagac 2280
 caaacagtat aaagcactca agaatacaca gttggaagtt actccaaaga atgagcacia 2340
 aacaatctta aagacactga aagatgagca gacaagaaaa cttgccattt tggcagagca 2400
 gtatgaacag agtataaatg aaatgatggc ctctcaagcg ttacggctag atgaggctca 2460
 agaagcagaa tgccaggcct tgaggctaca gctccagcag gaaatggagc tgctcaacgc 2520
 ctaccagagc aaaatcaaga tgcaaacaga ggcacaacat gaacgtgagc tccagaagct 2580
 agagcagaga gtgtctctgc gcagagcaca ccttgagcag aagattgaag aggagctggc 2640
 tgcccttcag aaggaacgca gcgagagaat aaagaacctt ttggaaaggc aagagcgaga 2700
 gattgaaact tttgacatgg agagcctcag aatgggattt gggaaatttg ttacattaga 2760
 ttttctaag gaggactaca gatgagatta aattttttgc catttac 2807

<210> 10

<211> 4620

<212> DNA

<213> Homo sapiens

<400> 10

gagccgtgat tgtgccacta cactccagcc ctgacctctt acaccgaagc agtcctcata 60
 cgtcagcctc ccaaagtgtt gggattacag atgaaccaag gatcgggata gcagtataaa 120
 attagaatca agacagctga ctgctcagca ggatgccatc aactaacaga gcaggcagcc 180
 tgaaggaccc tgaaattgca gagctottct tcaaagaaga tccagagaag ctcttcacag 240
 atctcagaga aattggccat ggaagctttg gagcagtgtt ttttgacga gatgtgcgtt 300
 ccaatgaagt ggtggccatc aagaaaatgt cttatagtgg aaagcagtct actgagaaat 360
 ggcaggatat tattaaggaa gtcaagtttc tacaagaat aaaacatccc aacagtatag 420
 aatacaaagg ctgttattta cgtgaacaca cagcatggct tgtaatggaa tattgtttag 480
 gatctgcttc ggatttacta gaagttcaca aaaagccatt acaagaagt gaaatagcag 540
 caattacaca tgggtgctct cagggattag cctacttaca ttctcatact atgattcata 600
 gagatatcaa agcaggaaat atccttctga cagaaccagg ccagggtgaaa cttgctgact 660
 ttggctctgc ttccatggca tcacctgcca attcctttgt gggaacgccg tattggatgg 720
 ccccagaagt aatttttagc atggatgaag gacaatatga tggcaaagta gatgtgtggt 780
 ctcttggaat aacatgtatt gaactagcgg aaaggaagcc tcctttattt aatatgaatg 840
 caatgagtgc cttatatcac atagcccaaa atgaatcccc tacactacag tctaatgaat 900
 ggtctgatta ttttcgcaac tttgtagatt cttgcctcca gaaaatccct caagatcgac 960
 ctacatcaga ggaactttta aagcacatat ttgttcttcg ggagcgccct gaaaccgtgt 1020

taatagatct cattcagagg acaaaggatg cagtaagaga gctggacaat ctgcagtatc 1080
 gaaagatgaa gaaactcctt ttccaggagg cacataatgg accagcagta gaagcacagg 1140
 aagaagaaga ggaacaagat catggtgttg gccggacagg aacagttaat agtgttgga 1200
 gtaatcaatc cattcccagc atgtccatca gtgccagcag ccaaagcagt agtgttaaca 1260
 gtcttccaga tgtctcagat gacaagagtg agctagacat gatggaggga gaccacacag 1320
 tgatgtctaa cagttctgtt atccatttaa aaccagagga agaaaattac agagaagagg 1380
 gagatcctag aacaagagca tcagatccac aatctccacc ccaagtatct cgtcacaat 1440
 cacactatcg taatcgagaa cactttgcta ctatacggac agcatcactg gttacgaggc 1500
 aatgcaaga acatgagcag gactctgagc ttagagaaca aatgtctggc tataagcgaa 1560
 tgaggcgaca acatcaaaag caactgatga ctctggaaaa caagctaaag gctgagatgg 1620
 atgaacatcg cctcagatta gacaaagatc ttgaaactca gcgtaacaat tttgctgcag 1680
 aatggagaa acttatcaag aaacaccagg ctgctatgga gaaaggaggct aaagtgatgt 1740
 ccaatgaaga gaaaaaattt cagcaacata ttcaggccca acagaagaaa gaactgaata 1800
 gttttctcga gtcccagaaa agagagtata aacttcgaaa agagcagctt aaagaggagc 1860
 taaatgaaaa ccagagtacc cccaaaaaag aaaaacagga gtggctttca aagcagaagg 1920
 agaataatac gcatttccaa gcagaagaag aagctaacct tcttcgacgt caaagacaat 1980
 acctagagct ggaatgccgt cgcttcaaga gaagaatgtt acttgggcgt cataacttag 2040
 agcaggacct tgtcaggagg gagttaaca aaagacagac tcagaaggac ttagagcatg 2100
 ccatgctact ccgacagcat gaatctatgc aagaactgga gttccgccac ctcaacacaa 2160
 ttcagaagat gcgctgtgag ttgatcagat tacagcatca aactgagctc actaaccagc 2220
 tggaatataa taagcgaaga gaacgagaac taagacgaaa gcatgtcatg gaagtctgac 2280
 aacagcctaa gagtttgaag tctaaagaac tccaaataaa aaagcagttt caggatacct 2340
 gcaaaatcca aaccagacag tacaaagcat taagaaatca cctgctggag actacaccaa 2400
 agagtgaaca caaagctgtt ctgaaacggc tcaaggagga acagaccggg aaattagcta 2460
 tcttggctga gcagtatgat cacagcatta atgaaatgct ctccacacaa gccctgcgtt 2520
 tggatgaagc acaggaagca gagtgccagg ttttgaagat gcagctgcag caggaaactgg 2580
 agctgttgaa tgcgtatcag agcaaaatca agatgcaagc tgaggcacia catgatcgag 2640
 agcttcgcga gcttgaacag agggctctcc tccggagggc actcttagaa caaaagattg 2700
 aagaagagat gttggctttg cagaatgagc gcacagaacg aatacgaagc ctgttggaac 2760
 gtcaagccag agagattgaa gcttttgact ctgaaagcat gagactaggt tttagtaata 2820
 tggctcttcc taatctctcc cctgaggcat tcagccacag ctaccgggga gcttctggtt 2880

ggtcacacaa ccctactggg ggtccaggac ctactgggg tcatcccatg ggtggccac 2940
 cacaagcttg gggccatcca atgcaagggtg gacccagcc atgggggtcac ccttcagggc 3000
 caatgcaagg ggtacctcga ggtagcagta tgggagtcg caatagcccc caggctctga 3060
 ggcggacagc ttctggggga cggacagagc agggcatgag cagaagcacg agtgtcactt 3120
 cacaatatc caatgggtca cacatgtctt atacataact taataattga gagtggcaat 3180
 tccgctggag ctgtctgcca aaagaaactg cctacagaca tcatcacagc agcctcctca 3240
 cttgggtact acagtgtgga agctgagtg atatggtata ttttattcat ttttgtaaag 3300
 cgttctgttt tgtgtttact aattgggatg tcatagtact tggctgccgg gtttgtttgt 3360
 ttttggggaa attttgaaaa gtggagttga tattaataat aaatgtgtat gtgtgtacat 3420
 atatatacac acacatacac atatattatg catgtggtga aaagaattgg ctagataggg 3480
 gatttttctg aacactgcaa aaatagaacg tagcaaaatg gcttcagtta tcaacttttg 3540
 gtgtctgtat cctaagaagt ttctgaaaag atctaaagcc ttttatccc atatcccaa 3600
 ttcttatgag ccactcacag caggcagcat atgttgaaat aagttattac tggtagacac 3660
 ctgcattgcc tcaccagtgt atttattttgt tattaatgt atctgacttc tcagcctcat 3720
 ttggactaaa aaaagaaagc agaaatccat gaacacattg cttctcggcc ttttggctaa 3780
 gatcaagtgt agaaatccat gaacactaaa ggacttcatt gattttttca gagagtagaa 3840
 aacaacttag tttttctttt ttctgaatg cgtcataggc ttgtgagtga tttttgtcca 3900
 ttcaattgtg cttctttgt attatgataa gatgggggta cttaggaga tcacaagttg 3960
 tgtgaggatt gcattaacaa acctatgagc cttcaatggg gaagaccaga agggtagag 4020
 gggccctgaa agttcatatg gtgggtatgt cccgcagcag agtgaggaga tgaagcttac 4080
 gtgtcctgac gttttgttgc ttatactgtg atatctcatc ctagctaagc tctataatgc 4140
 ccgagacccc aaacagtact tttactttgt ttgtacaaa acaaagacat atagccaata 4200
 caaatcaaat gccggaggtg tttgatgcca tttttgcaa ttgccatcta ttgaaattct 4260
 cgtcacacta catagacata attgttatct ccttttggct tatgtgattt tctgtttaca 4320
 agtagaatag ccaattattt aaatgtttag ttgccacagt gaaccaggag tcaactgagc 4380
 aatgacttta ccagctgctg actaatcttc atcaccactg tagattttgc tgcagtgcga 4440
 ggtcctctat ttttaattgc tgttttcgtt gctgcagtac tttacaaact tctagttcgt 4500
 tgagacttag tgaccatttg gcatcaagtt aacatcacac aataggaaac accacttcca 4560
 caagtctcaa gcctcagtgc taaagtacta ctgaaaagga actaggaagt ttggccaatt 4620

<210> 11

<211> 4536

<212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)..(4536)
 <223> "n" is A, C, G, or T

```

<400> 11
tacagatgaa ccaaggatcg ggatagcagt ataaaattag aatcaagaca gctgactgct      60
cagcaggatg ccatcaacta acagagcagg cagcctgaag gaccctgaaa ttgcagagct      120
cttøttcaaa gaagatccag agaagctctt cacagatctc agagaaattg gccatggaag      180
ctttggagca gtgtattttg cagcagatgt gcgtaccaat gaagtgggtgg ccatcaagaa      240
aatgtcttat agtggaaagc agtctactga gaaatggcag gatattatta aggaagtcaa      300
gtttctacaa agaataaaac atccnaaca gtatagaata caaaggctgt tatttacgtg      360
aacacacagc atggcttgta atggaatatt gttaggatc tgcttcggat ttactagaag      420
ttcacaaaaa gccattacaa gaagtggaaa tagcagcaat tacacatggt gctcttcagg      480
gattagccta cttacattct catactatga ttcatagaga tatcaaagca ggaaatatcc      540
ttctgacaga accaggccag gtgaaacttg ctgactttgg ctctgcttcc atggcatcac      600
ctgccaattc ctttgtggga acgccgtatt ggatggcccc agaagtaatt ttagccatgg      660
atgaaggaca atatgatggc aaagtagatg tgtggctctct tggaataaca tgtattgaac      720
tagcggaaag gaagcctcct ttatttaata tgaatgcaat gagtgcctta tatcacatag      780
cccaaatga atcccctaca ctacagtcta atgaatggtc tgattatttt cgcaactttg      840
tagattcttg cctccagaaa atccctcaag atcgacctac atcagaggaa cttttaaaagc      900
acatatttgt tcttcgggag cgccctgaaa ccgtgttaat agatctcatt cagaggacaa      960
aggatgcagt aagagagctg gacaatctgc agtatcgaaa gatgaagaaa ctctttttcc      1020
aggaggcaca taatggacca gcagtagaag cacaggaaga agaagaggaa caagatcatg      1080
gtgttggccg gacaggaaca gttaatagtg ttggaagtaa tcaatccatt cccagcatgt      1140
ccatcagtgc cagcagccaa agcagtagtg ttaacagtct tccagatgtc tcagatgaca      1200
agagtgcagt agacatgatg gagggagacc acacagtgat gtctaacagt tctgttatcc      1260
atttaaaacc agaggaagaa aattacagag aagagggaga tcctagaaca agagcatcag      1320
atccacaatc tccaccccaa gtatctcgtc acaaatcaca ctatcgtaat cgagaacact      1380
ttgctactat acggacagca tcactgggta cgaggcaaat gcaagaacat gagcaggact      1440
ctgagcttag agaacaaatg tctggctata agcgaatgag gcgacaacat caaaagcaac      1500
tgatgactct ggaaaacaag ctaaaggctg agatggatga acatcgctc agattagaca      1560

```

aagatcttga aactcagcgt aacaattttg ctgcagaaat ggagaaactt atcaagaaac	1620
accaggctgc tatggagaaa gaggctaaag tgatgtccaa tgaagagaaa aaatttcagc	1680
aacatattca ggcccaacag aagaaagaac tgaatagttt tctcgagtcc cagaaaagag	1740
agtataaact tcgaaaagag cagcttaaag aggagctaaa tgaaaaccag agtaccacca	1800
aaaaagaaaa acaggagtgg ctttcaaagc agaaggagaa tatacagcat ttccaagcag	1860
aagaagaagc taaccttctt cgacgtcaaa gacaatacct agagctggaa tgccgtcgct	1920
tcaagagaag aatgttactt gggcgtcata acttagagca ggacctgtc agggaggagt	1980
taaacaaaag acagactcag aaggacttag agcatgccat gctactccga cagcatgaat	2040
ctatgcaaga actggagttc cgccacctca acacaattca gaagatgcgc tgtgagttga	2100
tcagattaca gcatcaaact gagctcacta accagctgga atataataag cgaagagAAC	2160
gagaactaag acgaaagcat gtcattggaag ttccgacaaca gcctaagagt ttgaagtcta	2220
aagaactcca aataaaaaag cagtttcagg atacctgcaa aatccaaacc agacagtaca	2280
aagcattaag aaatcacctg ctggagacta caccaaagag tgagcācaaa gctgttctga	2340
aacggctcaa ggaggaacag acccggaat tagctatctt ggctgagcag tatgatcaca	2400
gcattaatga aatgctctcc acacaagccc tgcgtttgga tgaagcacag gaagcagagt	2460
gccaggtttt gaagatgcag ctgcagcagg aactggagct gttgaatgcg tatcagagca	2520
aaatcaagat gcaagctgag gcacaacatg atcgagagct tcgcgagctt gaacagaggg	2580
tctccctccg gagggcactc ttagaacaaa agattgaaga agagatgttg gctttgcaga	2640
atgagcgcac agaacgaata cgaagcctgt tggaacgtca agccagagag attgaagctt	2700
ttgactctga aagcatgaga ctaggtttta gtaatatggt cctttctaat ctctccctg	2760
aggcattcag ccacagctac ccgggagctt ctggttggtc acacaacctt actgggggtc	2820
caggacctca ctgggtcat cccatgggtg gcccaccaca agcttggggc catccaatgc	2880
aagggtgacc ccagccatgg ggtcacctt cagggccaat gcaaggggta cctcgaggta	2940
gcagtatggg agtccgcaat agccccagg ctctgaggcg gacagcttct gggggacgga	3000
cagagcaggg catgagcaga agcacgagtg tcacttcaca aatatccaat gggtcacaca	3060
tgtcttatac ataacttaat aattgagagt ggcaattccg ctggagctgt ctgcaaaaag	3120
aaactgccta cagacatcat cacagcagcc tcctcacttg ggtactacag tgtggaagct	3180
gagtgcatat ggtatatttt attcattttt gtaaagcgtt ctgttttggtg tttactaatt	3240
gggatgtcat agtacttggc tgccgggttt gtttgttttt ggggaaattt tgaaaagtgg	3300
agttgatatt aaaaataaat gtgtatgtgt gtacatatat atacacacac atacacatat	3360
attatgcatg tggtgaaaag aattggctag ataggggatt tttctgaaca ctgcaaaaat	3420

agaacgtagc aaaatggctt cagttatcac ttttgggtgt ctgtatccta agaagtttct 3480
 gaaaagatct aaagcctttt tatcccatat cccaaattct tatgagccac tcacagcagg 3540
 cagcatatgt tgaaataagt tattactggg acacacctgc attgcctcac cagtgtattt 3600
 atttgttatt aaattgatct gacttctcag cctcatttgg actaaaaaaaa gaaagcagaa 3660
 atccatgaac acattgcttc tgggcctttt ggctaagatc aagtgtagaa atccatgaac 3720
 actaaaggac ttcatlgatt ttttcagaga gtagaaaaca acttagtttt tcttttttcc 3780
 tgaatgcgtc ataggcttgt gaggatgttt tgtccattca attgtgcctt ctttgtatta 3840
 tgataagatg ggggtactta aggagatcac aagttgtgtg aggattgcat taacaaacct 3900
 atgagccttc aatggggaag accagaaggg tgagaggggc cctgaaagt catatgggtg 3960
 gtatgtcccg cagcagagtg aggagatgaa gcttacgtgt cctgacgttt tgttgcttat 4020
 actgtgatat ctcatcctag ctaagctcta taatgccaa gaccccaaac agtactttta 4080
 ctttgtttgt acaaaaacaa agacatatag ccaatacaaa tcaaagccg gaggtgtttg 4140
 atgccatatt tgcaaattgc catctattga aattctcgtc acactacata gacataattg 4200
 ttatctcctt ttggcttatg tgattttctg tttacaagta gaatagccaa ttatttaaatt 4260
 gtttagttgc cacagtgaac caggagtcac tgagccaatg actttaccag ctgctgacta 4320
 atcttcatca ccaactgtaga ttttgctgca tgtgcaggtc ctctattttt aattgctgtt 4380
 ttcgttgctg cagtacttta caaacttcta gttcgttgag acttagtgac catttggcat 4440
 caagttaaca tcacacaata ggaaacacca cttccacaag tctcaagcct cagtgtctaaa 4500
 gtactactga aaaggaacta ggaagtttgg ccaatt 4536

<210> 12
 <211> 4535
 <212> DNA
 <213> Homo sapiens

<400> 12
 tacagatgaa ccaaggatcg ggatagcagt ataaaattag aatcaagaca gctgactgct 60
 cagcaggatg ccatcaacta acagagcagg cagcctgaag gaccctgaaa ttgcagagct 120
 cttcttcaaa gaagatccag agaagctctt cacagatctc agagaaattg gccatggaag 180
 ctttgaggca gtgtattttg cacgagatgt gcgtaccaat gaagtgggtg ccatcaagaa 240
 aatgtcttat agtggaagc agtctactga gaaatggcag gatattatta aggaagtcaa 300
 gtttctacaa agaataaaac atcccaacag tatagaatac aaaggctgtt atttacgtga 360
 acacacagca tggcttgtaa tgggaatattg tttaggatct gcttcggatt tactagaagt 420
 tcacaaaaag ccattacaag aagtggaaat agcagcaatt acacatgggtg ctcttcaggg 480
 attagcctac ttacattctc atactatgat tcatagagat atcaaagcag gaaatctct 540

tctgacagaa ccaggccagg tgaaacttgc tgacttttggc tctgcttcca tggcatcacc	600
tgccaattcc tttgtgggaa cgccgtattg gatggcccca gaagtaattt tagccatgga	660
tgaaggacaa tatgatggca aagtagatgt gtggtctctt ggaataacat gtattgaact	720
agcggaaaagg aagcctcctt tattttaatat gaatgcaatg agtgccttat atcacatagc	780
ccaaaatgaa tcccctacac tacagtctaa tgaatggtct gattattttt gcaactttgt	840
agattcttgc ctccagaaaa tccctcaaga tcgacctaca tcagaggaac ttttaaagca	900
catatttggt ctccgggagc gccctgaaac cgtgttaata gatctcattc agaggacaaa	960
ggatgcagta agagagctgg acaatctgca gtatcgaaaag atgaagaaac tcctttttcca	1020
ggaggcacat aatggaccag cagtagaagc acaggaagaa gaagaggaac aagatcatgg	1080
tgttggccgg acaggaacag ttaatagtgt tggaagtaat caatccattc ccagcatgtc	1140
catcagtgcc agcagccaaa gcagtagtgt taacagtctt ccagatgtct cagatgacaa	1200
gagtgaagta gacatgatgg agggagacca cacagtgatg tctaacagtt ctgttatcca	1260
tttaaaacca gaggaagaaa attacagaga agaggagat cctagaacaa gagcatcaga	1320
tccacaatct ccacccaag tatctcgtca caaatcacac tatcgtaatc gagaacactt	1380
tgctactata cggacagcat cactgggttac gaggcaaatg caagaacatg agcaggactc	1440
tgagcttaga gaacaaatgt ctggctataa gogaatgagg cgacaacatc aaaagcaact	1500
gatgactctg gaaaacaagc taaaggctga gatggatgaa catcgctca gattagacaa	1560
agatcttgaa actcagcgta acaattttgc tgcagaaatg gagaaactta tcaagaaaca	1620
ccaggctgcc atggagaaaag aggctaaagt gatgtccaat gaagagaaaa aatttcagca	1680
acatattcag gcccaacaga agaaagaact gaatagtttt ctcgagtccc agaaaagaga	1740
gtataaactt cgaaaagagc agcttaaaga ggagctaaat gaaaaccaga gtacccccaa	1800
aaaagaaaaa caggagtggc tttcaaagca gaaggagaat atacagcatt tccaagcaga	1860
agaagaagct aaccttcttc gacgtcaaag acaataccta gagctggaat gccgtcgctt	1920
caagagaaga atgttacttg ggcgtcataa cttagagcag gaccttgtca gggaggagtt	1980
aaacaaaaga cagactcaga aggacttaga gcatgccatg ctactccgac agcatgaatc	2040
tatgcaagaa ctggagtcc gccacctcaa cacaattcag aagatgcgct gtgagttgat	2100
cagattacag catcaaactg agctcactaa ccagctggaa tataataagc gaagagaacg	2160
agaactaaga cgaaagcatg tcatggaagt tcgacaacag cctaagagtt tgaagtetaa	2220
agaactccaa ataaaaaagc agtttcagga tacctgcaaa atccaaacca gacagtacaa	2280
agcattaaga aatcacctgc tggagactac accaaagagt gagcaciaag ctgttctgaa	2340
acggctcaag gaggaacaga cccggaaatt agctatcttg gctgagcagt atgatcacag	2400

cattaatgaa atgctctcca cacaagccct gcgtttggat gaagcacagg aagcagagtg 2460
 ccaggttttg aagatgcagc tgcagcagga actggagctg ttgaatgcgt atcagagcaa 2520
 aatcaagatg caagctgagg cacaacatga tgcagagctt cgcgagcttg aacagagggg 2580
 ctccctccgg agggcactct tagaacaaaa gattgaagaa gagatgttgg ctttgcagaa 2640
 tgagcgcaca gaacgaatac gaagcctgtt ggaacgtcaa gccagagaga ttgaagcttt 2700
 tgactctgaa agcatgagac taggttttag taatatgggtc ctttctaate tctccctga 2760
 ggcattcagc cacagctacc cgggagcttc tgggtgggtca cacaacccta ctgggggtcc 2820
 aggacctcac tggggtcac ccatgggtgg cccaccacaa gcttggggcc atccaatgca 2880
 aggtggaccc cagccatggg gtcacccttc agggccaatg caaggggtac ctgcaggtag 2940
 cagtatggga gtccgcaata gccccaggc tctgaggcgg acagcttctg ggggacggac 3000
 ggagcagggc atgagcagaa gcacgagtgt cacttcacaa atatccaatg ggtcacacat 3060
 gtcttatata taacttaata attgagagt gcaattccgc tggagctgtc tgccaaaaga 3120
 aactgcctac agacatcatc acagcagcct cctcacttgg gtactacagt gtggaagctg 3180
 agtgcataatg gtatatatta ttcatttttg taaagcgttc tgttttgtgt ttactaattg 3240
 ggatgtcata gtacttggct gccgggtttg tttgtttttg gggaaatttt gaaaagtggg 3300
 gttgatatta aaaataaatg tgtatgtgtg tacatatata tacacacaca tacacatata 3360
 ttatgcatgt ggtgaaaaga attggctaga taggggattt ttctgaacac tgcaaaaata 3420
 gaacgtagca aaatggcttc agttatcact tttgggtgtc tgtatcctaa gaagtgtctg 3480
 aaaagatcta aagccttttt atcccatatc ccaaattctt atgagccact cacagcaggc 3540
 agcatatgtt gaaataagtt attactggta cacacctgca ttgcctcacc agtgtattta 3600
 tttgttatta aattgatctg acttctcagc ctcatattgga ctaaaaaag aaagcagaaa 3660
 tccatgaaca cattgcttct cggccttttg gctaagatca agtgtagaaa tccatgaaca 3720
 cttaaaggact tcattgattt tttcagagag tagaaaacaa cttagttttt cttttttcct 3780
 gaatgcgtca taggttgtg agtgattttt gtccattcaa ttgtgccttc tttgtattat 3840
 gataagatgg ggggtactta ggagatcaca agttgtgtga ggattgcatt aacaaaccta 3900
 tgagccttca atggggaaga ccagaagggt gagagggggc ctgaaagtcc atatgggtgg 3960
 tatgtccgc agcagagtga ggagatgaag cttacgtgtc ctgacgtttt gttgcttata 4020
 ctgtgatatc tcactcctagc taagctctat aatgcccaag accccaaaca gtacttttac 4080
 tttgtttgta caaaaacaaa gacatatagc caatacaaat caaatgccgg aggtgtttga 4140
 tgccatattt gcaaattgcc atctattgaa attctcgtca cactacatag acataattgt 4200
 tatctccttt tggcttatgt gattttctgt ttacaagtag aatagccaat tatttaaatg 4260

ttttagttgcc acagtgaacc aggagtcact gagccaatga ctttaccagc tgctgactaa 4320
 tcttcatcac cactgtagat tttgctgcat gtgcaggctc tctattttta attgctgttt 4380
 tcgttgctgc agtactttac aaactttctag ttcgttgaga cttagtgacc atttggcatc 4440
 aagttaacat cacacaatag gaaacaccac ttccacaagt ctcaagcctc agtgctaaag 4500
 tactactgaa aaggaactag gaagtttggc caatt 4535

<210> 13
 <211> 3003
 <212> DNA
 <213> Homo sapiens

<400> 13
 atgccatcaa ctaacagagc aggcagtctt aaggaccctg aaattgcaga gctcttcttc 60
 aaagaagatc cagagaagct cttcacagat ctccagagaaa ttggccatgg aagcttttga 120
 gcagtgtatt ttgcacgaga tgtgcgtacc aatgaagtgg tggccatcaa gaaaatgtct 180
 tatagtggaa agcagtctac tgagaaatgg caggatatta ttaaggaagt caagtttcta 240
 caaagaataa aacatcccaa cagtatagaa tacaaggct gttatttacg tgaacacaca 300
 gcatggcttg taatggaata ttgttttagga tctgcttcgg atttactaga agttcacaaa 360
 aagccattac aagaagtgg aatagcagca attacacatg gtgctcttca gggattagcc 420
 tacttacatt ctccatactat gattcataga gatatcaaag caggaaatat ccttctgaca 480
 gaaccaggcc aggtgaaact tgctgacttt ggctctgctt ccatggcatc acctgccaat 540
 tcctttgtgg gaacgccgta ttggatggcc ccagaagtaa ttttagccat ggatgaagga 600
 caatatgatg gcaaagtaga tgtgtggtct cttggaataa catgtattga actagcggaa 660
 aggaagcctc ctttatttaa tatgaatgca atgagtgcct tatatcacat agcccaaat 720
 gaatcccta cactacagtc taatgaatgg tctgattatt ttgcgaactc tgtagattct 780
 tgctccaga aaatccctca agatcgacct acatcagagg aacttttaaa gcacatattt 840
 gttcttcggg agcgccctga aaccgtgtta atagatctca ttcagaggac aaaggatgca 900
 gtaagagagc tggacaatct gcagtatcga aagatgaaga aactcctttt ccaggaggca 960
 cataatggac cagcagtaga agcacaggaa gaagaagagg aacaagatca tgggtgttggc 1020
 cggacaggaa cagttaatag tgttggaagt aatcaatcca ttcccagcat gtccatcagt 1080
 gccagcagcc aaagcagtag tgttaacagt cttccagatg tctcagatga caagagttag 1140
 ctagacatga tggagggaga ccacacagtg atgtctaaca gttctgttat ccatttaaaa 1200
 ccagaggaag aaaattacag agaagaggga gatcctagaa caagagcatc agatccacaa 1260
 tctccacccc aagtatctcg tcacaaatca cactatcgta atcgagaaca ctttgcact 1320

atacggacag catcactggt tacgaggcaa atgcaagaac atgagcagga ctctgagctt 1380
 agagaacaaa tgtctggcta taagcgaatg aggcgacaac atcaaaagca actgatgact 1440
 ctggaaaaaca agctaaaggc tgagatggat gaacatcgcc tcagattaga caaagatctt 1500
 gaaactcagc gtaacaatth tgctgcagaa atggagaaac ttatcaagaa acaccaggct 1560
 gctatggaga aagaggctaa agtgaatgcc aatgaagaga aaaaatttca gcaacatatt 1620
 caggcccaac agaagaaaga actgaatagt tttctcgagt cccagaaaag agagtataaa 1680
 cttcgaaaag agcagcttaa agaggagcta aatgaaaacc agagtacccc caaaaaagaa 1740
 aaacaggagt ggctttcaaa gcagaaggag aatatacagc atttccaagc agaagaagaa 1800
 gctaaccttc ttgcagctca aagacaatac ctagagctgg aatgccgtcg cttcaagaga 1860
 agaatgttac ttgggcgtca taacttagag caggaccttg tcaggaggga gttaaataaa 1920
 agacagactc agaaggactt agagcatgcc atgctactcc gacagcatga atctatgcaa 1980
 gaactggagt tccgccacct caacacaatt cagaagatgc gctgtgagtt gatcagatta 2040
 cagcatcaaa ctgagctcac taaccagctg gaatataata agcgaagaga acgagaacta 2100
 agacgaaagc atgtcatgga agttcgacaa cagcctaaga gtttgaagtc taaagaactc 2160
 caaataaaaa agcagtttca ggatacctgc aaaatccaaa ccagacagta caaagcatta 2220
 agaaatcacc tggtggagac tacaccaaag agtgagcaca aagctgttct gaaacggctc 2280
 aaggaggaac agaccgggaa attagctatc ttggctgagc agtatgatca cagcattaat 2340
 gaaatgctct ccacacaagc cctgcgtttg gatgaagcac aggaagcaga gtgccagggt 2400
 ttgaagatgc agctgcagca ggaactggag ctggttgaatg cgtatcagag caaaatcaag 2460
 atgcaagctg aggcacaaca tgatcgagag cttcgcgagc ttgaacagag ggtctccctc 2520
 cggagggcac tcttagaaca aaagattgaa gaagagatgt tggctttgca gaatgagtgc 2580
 acagaacgaa tacgaagcct gttggaacgt caagccagag agattgaagc ttttgactct 2640
 gaaagcatga gactaggtht tagtaatatg gtcctttcta atctctcccc tgaggcattc 2700
 agccacagct acccgggagc ttctgggttg tcacacaacc ctactggggg tccaggacct 2760
 cactggggtc atcccattggg tggcccaacca caagcttggg gccatccaat gcaagggtga 2820
 cccagccat ggggtcacc ttcagggccca atgcaagggg tacctcgagg tagcagtatg 2880
 ggagtccgca atagcccca ggctctgagg cggacagctt ctgggggacg gacagagcag 2940
 ggcattgagca gaagcacgag tgtcacttca caaatatcca atgggtcaca catgtcttat 3000
 aca 3003

<210> 14
 <211> 3048
 <212> DNA

<213> Homo sapiens

<400> 14

tgctcagcag gatgccatca actaacagag caggcagcct gaaggaccct gaaattgcag	60
agctcttctt caaagaagat ccagagaagc tcttcacaga tctcagagaa attggccatg	120
gaagcttttg agcagtgtat tttgcacgag atgtgcgtac caatgaagtg gtggccatca	180
agaaaatgtc ttatagtgga aagcagtcta ctgagaaatg gcaggatatt attaaggaag	240
tcaagtttct acaaagaata aaacatccca acagtataga atacaaaggc tgttattttac	300
gtgaacacac agcatggcct gtaatggaat attgttttagg atctgcttcg gatttactag	360
aagttcacaa aaagccatta caagaagtgg aaatagcagc aattacacat ggtgctcttc	420
agggattagc ctactttacat tctcatacta tgattcatag agatatcaaa gcaggaaata	480
tccttctgac agaaccaggc caggtgaaac ttgctgactt tggctctgct tccatggcat	540
cacctgccaa ttccttttggt ggaacgccgt attggatggc cccagaagta atttttagcca	600
tggatgaagg acaatatgat ggcaaagtag atgtgtgggc tcttggaata acatgtattg	660
aactagcgga aaggaagcct cctttattta atatgaatgc aatgagtgcc ttatatcaca	720
tagcccaaaa tgaatcccct acactacagt ctaatgaatg gtctgattat tttcgcaact	780
ttgtagattc ttgcctccag aaaatccctc aagatcgacc tacatcagag gaacttttaa	840
agcacatatt tgttcttcgg gagcgccctg aaaccgtgtt aatagatctc attcagagga	900
caaaggatgc agtaagagag ctggacaatc tgcagtatcg aaagatgaag aaactccttt	960
tccaggaggc acataatgga ccagcagtag aagcacagga agaagaagag gaacaagatc	1020
atggtgttgg ccggacagga acagttaata gtgttggaag taatcaatcc attccagca	1080
tgtccatcag tgccagcagc caaagcagta gtgttaacag tcttcagat gtctcagatg	1140
acaagagtga gctagacatg atggagggag accacacagt gatgtctaac agttctgtta	1200
tccattttaa accagaggaa gaaaattaca gagaagaggg agatcctaga acaagagcat	1260
cagatccaca atctccaccc caagtatctc gtcacaaatc acactatcgt aatcgagAAC	1320
actttgtac tatacgga gcatcactgg ttacgaggca aatgcaagaa catgagcagg	1380
actctgagct tagagaacaa atgtctggct ataagcgaat gaggcgacaa catcaaaagc	1440
aactgatgac tctggaaaac aagctaaagg ctgagatgga tgaacatcgc ctcagattag	1500
acaaagatct tgaaactcag cgtaacaatt ttgctgcaga aatggagaaa cttatcaaga	1560
aacaccaggc tgctatggag aaagaggcta aagtgatgtc caatgaagag aaaaaatttc	1620
agcaacatat tcaggcccaa cagaagaaag aactgaatag ttttctcgag tcccagaaaa	1680
gagagtataa acttcgaaaa gagcagctta aagaggagct aaatgaaaac cagagtaccc	1740
ccaaaaaaga aaaacaggag tggctttcaa agcagaagga gaatatacag catttccaag	1800

cagaagaaga agctaaccctt cttcgacgtc aaagacaata cctagagctg gaatgccgtc 1860
 gcttcaagag aagaatgtta cttggggcgtc ataacttaga gcaggacctt gtcagggagg 1920
 agttaaacia aagacagact cagaaggact tagagcatgc catgctactc cgacagcatg 1980
 aatctatgca agaactggag ttccgccacc tcaacacaat tcagaagatg cgctgtgagt 2040
 tgatcagatt acagcatcaa actgagctca ctaaccagct ggaatataat aagcgaagag 2100
 aacgagaact aagacgaaag catgtcatgg aagtctgaca acagcctaag agtttgaagt 2160
 ctaaagaact ccaaataaaa aagcagtttc aggatacctg caaaatccaa accagacagt 2220
 acaaagcatt aagaaatcac ctgctggaga ctacaccaaa gagtgcagcac aaagctgttc 2280
 tgaaacggct caaggaggaa cagaccggga aattagctat cttggctgag cagtatgatc 2340
 acagcattaä tgaaatgtc tccacacaag ccctgcgttt ggatgaagca caggaagcag 2400
 agtgccagggt tttgaagatg cagctgcagc aggaactgga gctgttgaat gcgtatcaga 2460
 gcaaaatcaa gatgcaagct gaggcacaac atgatcgaga gcttcgcgag cttgaacaga 2520
 gggctctccct ccggagggca ctcttagaac aaaagattga agaagagatg ttggctttgc 2580
 agaatgagcg cacagaacga atacgaagcc tgttggaacg tcaagccaga gagattgaag 2640
 cttttgactc tgaaagcatg agactagggt ttagtaatat ggtcctttct aatctctccc 2700
 ctgaggcatt cagccacagc taccggggag cttctgggtg gtcacacaac cctactgggg 2760
 gtccaggacc tcaactgggggt catcccatgg gtggcccacc acaagcttgg ggccatccaa 2820
 tgcaagggtg accccagcca tggggtcacc cttcagggcc aatgcaagggt gtacctcgag 2880
 gtagcagtat gggagtccgc aatagcccc aggctctgag gcggacagct tctgggggac 2940
 ggacagagca gggcatgagc agaagcacga gtgtcacttc acaaatatcc aatgggtcac 3000
 acatgtctta tacataactt aataattgag agtggcaatt ccgctgga 3048

<210> 15
 <211> 3148
 <212> DNA
 <213> Homo sapiens

<400> 15
 ggcacgaggg tggcgccggg cggcgggggtc ctgcgtggag agtgggacgc aacgccgaga 60
 ccgcgagcag aggtgcgca cagccggatc cggcactcag cgaccggacc caaggatccg 120
 ccggggaaca agccacagga gagcgactca ggaacaagtg tgggagagga agcggcgggcg 180
 gcggcgccgg gccgggggt ggtgacagca ggtctgaggt tgcatacataa atacaaagga 240
 ctgaagttat aaaagagaaa agagaagttt gctgctaaaa tgaatctgag caatatggaa 300
 tattttgtgc cacacacaaa aaggctactga agattttacc cccaaaaaaa attgtcaatg 360

agaaataaag ctaactgata tcaaaaagca gagcctgctc tactggccat catgcgtaaa 420
 ggggtgctga aggaccaga gattgccgat ctattctaca aagatgatcc tgaggaactt 480
 tttattgggt tgcataaaat tggacatgga agttttggag cagtttattt tgctacaaat 540
 gctcacacca gtgaggtggt ggcaattaag aagatgtcct atagtgggaa gcagaccat 600
 gagaaatggc aagatattct taaggaagt aaatttttac gacaattgaa gcatcctaata 660
 actattgagt acaaaggctg ttacttgaaa gaacacactg cttggttggt gatggaatat 720
 tgcttaggct cagcctctga tttattagaa gtccataaaa aaccacttca ggaagtggag 780
 atcgtgccca ttactcatgg agccttgcat ggactagcct acctacattc tcatgcattg 840
 attcataggg atattaaagc aggaatatt. cttctaacag agccagggtca ggtaaaacta 900
 gctgattttg gatctgcttc aatggcttct cctgccaaact cttcgtggg cacaccttac 960
 tggatggctc cagaggtgat cttagctatg gatgaaggac agtatgatgg gaaagtgtgat 1020
 atttggtcac ttggcatcac ttgtattgaa ttggcggaac ggaagccgcc ctttttcaac 1080
 atgaatgcaa tgagtgcctt ataccacatt gccagaaatg actcccaac gttacagtct 1140
 aatgaatgga cagactcctt taggagattt gttgattact gcttgagaa aatacctcag 1200
 gaaaggccaa catcagcaga actattaagg catgactttg ttcgacgaga ccggccacta 1260
 cgtgtcctca ttgacctcat acagaggaca aaagatgcag ttcgtgagct agataacct 1320
 cagtaccgaa aaatgaaaaa aatacttttc caagagacac ggaatggacc cttgaatgag 1380
 tcacaggagg atgaggaaga cagtgaacat ggaaccagcc tgaacaggga aatggacagc 1440
 ctgggcagca accattccat tccaagcatg tccgtgagca caggcagcca gagcagcagt 1500
 gtgaacagca tgcaggaagt catggacgag agcagttccg aacttgatcat gatgcacgat 1560
 gacgaaagca caatcaattc cagctcctcc gtcgtgcata agaaagatca tgtattcata 1620
 agggatgagg cgggccacgg cgatcccagg cctgagccgc ggcctacca gtcagttcag 1680
 agccaggccc tccactaccg gaacagagag cgctttgccca cgatcaaata agcatctttg 1740
 gttacacgac agatccatga gcatgagcag gagaacgagt tgcgggaaca gatgtcaggt 1800
 tataagcgga tgcggcgcca gcaccagaag cagctgatcg ccctggagaa caagctgaag 1860
 gctgagatgg acgagcaccg cctcaagcta cagaaggagg tggagacgca tgccaacaac 1920
 tcgtccatcg agctggagaa gctggccaag aagcaagtgg ctatcataga aaaggaggca 1980
 aaggtagctg cagcagatga gaagaagttc cagcaacaga tcttggccca gcagaagaaa 2040
 gatttgacaa ctttcttaga aagtcagaag aagcagtata agatttgtaa gaaaaaata 2100
 aaagaggaaa tgaatgagga ccatagcaca ccaagaaaag agaagcaaga gcggatctcc 2160
 aaacataaag agaacttgca gcacacacag gctgaagagg aagccacct tctactcaa 2220

cagagactgt actacgacaa aaattgtcgt ttcttcaagc ggaaaataat gatcaagcgg 2280
 cactgaggtgg agcagcagaa cattcggggag gaactaaata aaaagaggac ccagaaggag 2340
 atggagcatg ccatgctaata cgggcacgac gagtccaccc gagagctaga gtacaggcag 2400
 ctgcacacgt tacagaagct acgcatggat ctgatccgtt tacagcacca gacggaactg 2460
 gaaaaccagc tggagtacaa taagaggcga gaaagagaac tgcacagaaa gcatgtcatg 2520
 gaacttcggc aacagccaaa aaacttaaag gccatggaaa tgcaaattaa aaaacagttt 2580
 caggacactt gcaaagtaca gaccaaacag tataaagcac tcaagaatca ccagttggaa 2640
 gttactccaa agaattgagca caaaacaatc ttaaagacac tgaaagatga gcagacaaga 2700
 aaacttgcca ttttggcaga gcagtatgaa cagagtataa atgaaatgat ggcctctcaa 2760
 gcgttacggc tagatgaggc tcaagaagca gaatgccagg ccttgaggct acagctccag 2820
 caggaaatgg agctgctcaa cgcctaccag agcaaaatca agatgcaaac agaggcacia 2880
 catgaacgtg agctccagaa gctagagcag agagtgtctc tgcgcagagc acaccttgag 2940
 cagaagattg aagaggagct ggctgccctt cagaaggaac gcagcgagag aataaagaac 3000
 ctattggaaa ggcaagagcg agagattgaa acttttgaca tggagagcct cagaatggga 3060
 tttgggaatt tggttacatt agattttcct aaggaggact acagatgaga ttaaattttt 3120
 tgccatttac aaaaaaaaaa aaaaaaaaaa 3148

<210> 16
 <211> 1049
 <212> PRT
 <213> Homo sapiens

<400> 16

Met Pro Ala Gly Gly Arg Ala Gly Ser Leu Lys Asp Pro Asp Val Ala
 1 5 10 15

Glu Leu Phe Phe Lys Asp Asp Pro Glu Lys Leu Phe Ser Asp Leu Arg
 20 25 30

Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe Ala Arg Asp Val
 35 40 45

Arg Asn Ser Glu Val Val Ala Ile Lys Lys Met Ser Tyr Ser Gly Lys
 50 55 60

Gln Ser Asn Glu Lys Trp Gln Asp Ile Ile Lys Glu Val Arg Phe Leu
 65 70 75 80

Gln Lys Leu Arg His Pro Asn Thr Ile Gln Tyr Arg Gly Cys Tyr Leu
 85 90 95

Arg Glu His Thr Ala Trp Leu Val Met Glu Tyr Cys Leu Gly Ser Ala
 100 105 110

Ser Asp Leu Leu Glu Val His Lys Lys Pro Leu Gln Glu Val Glu Ile
 115 120 125

Ala Ala Val Thr His Gly Ala Leu Gln Gly Leu Ala Tyr Leu His Ser
 130 135 140

His Asn Met Ile His Arg Asp Val Lys Ala Gly Asn Ile Leu Leu Ser
 145 150 155 160

Glu Pro Gly Leu Val Lys Leu Gly Asp Phe Gly Ser Ala Ser Ile Met
 165 170 175

Ala Pro Ala Asn Ser Phe Val Gly Thr Pro Tyr Trp Met Ala Pro Glu
 180 185 190

Val Ile Leu Ala Met Asp Glu Gly Gln Tyr Asp Gly Lys Val Asp Val
 195 200 205

Trp Ser Leu Gly Ile Thr Cys Ile Glu Leu Ala Glu Arg Lys Pro Pro
 210 215 220

Leu Phe Asn Met Asn Ala Met Ser Ala Leu Tyr His Ile Ala Gln Asn
 225 230 235 240

Glu Ser Pro Val Leu Gln Ser Gly His Trp Ser Glu Tyr Phe Arg Asn
 245 250 255

Phe Val Asp Ser Cys Leu Gln Lys Ile Pro Gln Asp Arg Pro Thr Ser
 260 265 270

Glu Val Leu Leu Lys His Arg Phe Val Leu Arg Glu Arg Pro Pro Thr
 275 280 285

Val Ile Met Asp Leu Ile Gln Arg Thr Lys Asp Ala Val Arg Glu Leu
 290 295 300

Asp Asn Leu Gln Tyr Arg Lys Met Lys Lys Ile Leu Phe Gln Glu Ala
 305 310 315 320

Pro Asn Gly Pro Gly Ala Glu Ala Pro Glu Glu Glu Glu Glu Ala Glu
 325 330 335

Pro Tyr Met His Arg Ala Gly Thr Leu Thr Ser Leu Glu Ser Ser His
 340 345 350

Ser Val Pro Ser Met Ser Ile Ser Ala Ser Ser Gln Ser Ser Ser Val
 355 360 365

Asn Ser Leu Ala Asp Ala Ser Asp Asn Glu Glu Glu Glu Glu Glu
 370 375 380

Glu Glu Glu Glu Glu Glu Glu Glu Gly Pro Glu Ala Arg Glu Met Ala
 385 390 395 400

Met Met Gln Glu Gly Glu His Thr Val Thr Ser His Ser Ser Ile Ile
 405 410 415

His Arg Leu Pro Gly Ser Asp Asn Leu Tyr Asp Asp Pro Tyr Gln Pro
 420 425 430

Glu Ile Thr Pro Ser Pro Leu Gln Pro Pro Ala Ala Pro Ala Pro Thr
 435 440 445

Ser Thr Thr Ser Ser Ala Arg Arg Arg Ala Tyr Cys Arg Asn Arg Asp
 450 455 460

His Phe Ala Thr Ile Arg Thr Ala Ser Leu Val Ser Arg Gln Ile Gln
 465 470 475 480

Glu His Glu Gln Asp Ser Ala Leu Arg Glu Gln Leu Ser Gly Tyr Lys
 485 490 495

Arg Met Arg Arg Gln His Gln Lys Gln Leu Leu Ala Leu Glu Ser Arg
 500 505 510

Leu Arg Gly Glu Arg Glu Glu His Ser Ala Arg Leu Gln Arg Glu Leu
 515 520 525

Glu Ala Gln Arg Ala Gly Phe Gly Ala Glu Ala Glu Lys Leu Ala Arg
 530 535 540

Arg His Gln Ala Ile Gly Glu Lys Glu Ala Arg Ala Ala Gln Ala Glu
 545 550 555 560

Glu Arg Lys Phe Gln Gln His Ile Leu Gly Gln Gln Lys Lys Glu Leu
 565 570 575

Ala Ala Leu Leu Glu Ala Gln Lys Arg Thr Tyr Lys Leu Arg Lys Glu
 580 585 590

Gln Leu Lys Glu Glu Leu Gln Glu Asn Pro Ser Thr Pro Lys Arg Glu
595 600 605

Lys Ala Glu Trp Leu Leu Arg Gln Lys Glu Gln Leu Gln Gln Cys Gln
610 615 620

Ala Glu Glu Glu Ala Gly Leu Leu Arg Arg Gln Arg Gln Tyr Phe Glu
625 630 635 640

Leu Gln Cys Arg Gln Tyr Lys Arg Lys Met Leu Leu Ala Arg His Ser
645 650 655

Leu Asp Gln Asp Leu Leu Arg Glu Asp Leu Asn Lys Lys Gln Thr Gln
660 665 670

Lys Asp Leu Glu Cys Ala Leu Leu Leu Arg Gln His Glu Ala Thr Arg
675 680 685

Glu Leu Glu Leu Arg Gln Leu Gln Ala Val Gln Arg Thr Arg Ala Glu
690 695 700

Leu Thr Arg Leu Gln His Gln Thr Glu Leu Gly Asn Gln Leu Glu Tyr
705 710 715 720

Asn Lys Arg Arg Glu Gln Glu Leu Arg Gln Lys His Ala Ala Gln Val
725 730 735

Arg Gln Gln Pro Lys Ser Leu Lys Ser Lys Glu Leu Gln Ile Lys Lys
740 745 750

Gln Phe Gln Glu Thr Cys Lys Ile Gln Thr Arg Gln Tyr Lys Ala Leu
755 760 765

Arg Ala His Leu Leu Glu Thr Thr Pro Lys Ala Gln His Lys Ser Leu
770 775 780

Leu Lys Arg Leu Lys Glu Glu Gln Thr Arg Lys Leu Ala Ile Leu Ala
785 790 795 800

Glu Gln Tyr Asp Gln Ser Ile Ser Glu Met Leu Ser Ser Gln Ala Leu
805 810 815

Arg Leu Asp Glu Thr Gln Glu Ala Glu Phe Gln Ala Leu Arg Gln Gln
820 825 830

Leu Gln Gln Glu Leu Glu Leu Leu Asn Ala Tyr Gln Ser Lys Ile Lys
835 840 845

Ile Arg Thr Glu Ser Gln His Glu Arg Glu Leu Arg Glu Leu Glu Gln
850 855 860

Arg Val Ala Leu Arg Arg Ala Leu Leu Glu Gln Arg Val Glu Glu Glu
865 870 875 880

Leu Leu Ala Leu Gln Thr Gly Arg Ser Glu Arg Ile Arg Ser Leu Leu
885 890 895

Glu Arg Gln Ala Arg Glu Ile Glu Ala Phe Asp Ala Glu Ser Met Arg
900 905 910

Leu Gly Phe Ser Ser Met Ala Leu Gly Gly Ile Pro Ala Glu Ala Ala
915 920 925

Ala Gln Gly Tyr Pro Ala Pro Pro Pro Ala Pro Ala Trp Pro Ser Arg
930 935 940

Pro Val Pro Arg Ser Gly Ala His Trp Ser His Gly Pro Pro Pro Pro
945 950 955 960

Gly Met Pro Pro Pro Ala Trp Arg Gln Pro Ser Leu Leu Ala Pro Pro
965 970 975

Gly Pro Pro Asn Trp Leu Gly Pro Pro Thr Gln Ser Gly Thr Pro Arg
980 985 990

Gly Gly Ala Leu Leu Leu Leu Arg Asn Ser Pro Gln Pro Leu Arg Arg
995 1000 1005

Ala Ala Ser Gly Gly Ser Gly Ser Glu Asn Val Gly Pro Pro Ala
1010 1015 1020

Ala Ala Val Pro Gly Pro Leu Ser Arg Ser Thr Ser Val Ala Ser
1025 1030 1035

His Ile Leu Asn Gly Ser Ser His Phe Tyr Ser
1040 1045

<210> 17

<211> 898

<212> PRT

<213> Homo sapiens

<400> 17

Met Arg Lys Gly Val Leu Lys Asp Pro Glu Ile Ala Asp Leu Phe Tyr
 1 5 10 15
 Lys Asp Asp Pro Glu Glu Leu Phe Ile Gly Leu His Glu Ile Gly His
 20 25 30
 Gly Ser Phe Gly Ala Val Tyr Phe Ala Thr Asn Ala His Thr Ser Glu
 35 40 45
 Val Val Ala Ile Lys Lys Met Ser Tyr Ser Gly Lys Gln Thr His Glu
 50 55 60
 Lys Trp Gln Asp Ile Leu Lys Glu Val Lys Phe Leu Arg Gln Leu Lys
 65 70 75 80
 His Pro Asn Thr Ile Glu Tyr Lys Gly Cys Tyr Leu Lys Glu His Thr
 85 90 95
 Ala Trp Leu Val Met Glu Tyr Cys Leu Gly Ser Ala Ser Asp Leu Leu
 100 105 110
 Glu Val His Lys Lys Pro Leu Gln Glu Val Glu Ile Ala Ala Ile Thr
 115 120 125
 His Gly Ala Leu His Gly Leu Ala Tyr Leu His Ser His Ala Leu Ile
 130 135 140
 His Arg Asp Ile Lys Ala Gly Asn Ile Leu Leu Thr Glu Pro Gly Gln
 145 150 155 160
 Val Lys Leu Ala Asp Phe Gly Ser Ala Ser Met Ala Ser Pro Ala Asn
 165 170 175
 Ser Phe Val Gly Thr Pro Tyr Trp Met Ala Pro Glu Val Ile Leu Ala
 180 185 190
 Met Asp Glu Gly Gln Tyr Asp Gly Lys Val Asp Ile Trp Ser Leu Gly
 195 200 205
 Ile Thr Cys Ile Glu Leu Ala Glu Arg Lys Pro Pro Leu Phe Asn Met
 210 215 220
 Asn Ala Met Ser Ala Leu Tyr His Ile Ala Gln Asn Asp Ser Pro Thr
 225 230 235 240
 Leu Gln Ser Asn Glu Trp Thr Asp Ser Phe Arg Arg Phe Val Asp Tyr

38

Val Glu Thr His Ala Asn Asn Ser Ser Ile Glu Leu Glu Lys Leu Ala
500 505 510

Lys Lys Gln Val Ala Ile Ile Glu Lys Glu Ala Lys Val Ala Ala Ala
515 520 525

Asp Glu Lys Lys Phe Gln Gln Gln Ile Leu Ala Gln Gln Lys Lys Asp
530 535 540

Leu Thr Thr Phe Leu Glu Ser Gln Lys Lys Gln Tyr Lys Ile Cys Lys
545 550 555 560

Glu Lys Ile Lys Glu Glu Met Asn Glu Asp His Ser Thr Pro Lys Lys
565 570 575

Glu Lys Gln Glu Arg Ile Ser Lys His Lys Glu Asn Leu Gln His Thr
580 585 590

Gln Ala Glu Glu Glu Ala His Leu Leu Thr Gln Gln Arg Leu Tyr Tyr
595 600 605

Asp Lys Asn Cys Arg Phe Phe Lys Arg Lys Ile Met Ile Lys Arg His
610 615 620

Glu Val Glu Gln Gln Asn Ile Arg Glu Glu Leu Asn Lys Lys Arg Thr
625 630 635 640

Gln Lys Glu Met Glu His Ala Met Leu Ile Arg His Asp Glu Ser Thr
645 650 655

Arg Glu Leu Glu Tyr Arg Gln Leu His Thr Leu Gln Lys Leu Arg Met
660 665 670

Asp Leu Ile Arg Leu Gln His Gln Thr Glu Leu Glu Asn Gln Leu Glu
675 680 685

Tyr Asn Lys Arg Arg Glu Arg Glu Leu His Arg Lys His Val Met Glu
690 695 700

Leu Arg Gln Gln Pro Lys Asn Leu Lys Ala Met Glu Met Gln Ile Lys
705 710 715 720

Lys Gln Phe Gln Asp Thr Cys Lys Val Gln Thr Lys Gln Tyr Lys Ala
725 730 735

Leu Lys Asn His Gln Leu Glu Val Thr Pro Lys Asn Glu His Lys Thr

```
<210> 18
<211> 1005
<212> PRT
<213> Homo sapiens

<400> 18
```

Leu	Leu	Ser	Arg	Met	Pro	Ser	Thr	Asn	Arg	Ala	Gly	Ser	Leu	Lys	Asp
1				5					10					15	
Pro	Glu	Ile	Ala	Glu	Leu	Phe	Phe	Lys	Glu	Asp	Pro	Glu	Lys	Leu	Phe
			20					25					30		
Thr	Asp	Leu	Arg	Glu	Ile	Gly	His	Gly	Ser	Phe	Gly	Ala	Val	Tyr	Phe
		35					40					45			

Ala Arg Asp Val Arg Thr Asn Glu Val Val Ala Ile Lys Lys Met Ser
50 55 60

Tyr Ser Gly Lys Gln Ser Thr Glu Lys Trp Gln Asp Ile Ile Lys Glu
65 70 75 80

Val Lys Phe Leu Gln Arg Ile Lys His Pro Asn Ser Ile Glu Tyr Lys
85 90 95

Gly Cys Tyr Leu Arg Glu His Thr Ala Trp Leu Val Met Glu Tyr Cys
100 105 110

Leu Gly Ser Ala Ser Asp Leu Leu Glu Val His Lys Lys Pro Leu Gln
115 120 125

Glu Val Glu Ile Ala Ala Ile Thr His Gly Ala Leu Gln Gly Leu Ala
130 135 140

Tyr Leu His Ser His Thr Met Ile His Arg Asp Ile Lys Ala Gly Asn
145 150 155 160

Ile Leu Leu Thr Glu Pro Gly Gln Val Lys Leu Ala Asp Phe Gly Ser
165 170 175

Ala Ser Met Ala Ser Pro Ala Asn Ser Phe Val Gly Thr Pro Tyr Trp
180 185 190

Met Ala Pro Glu Val Ile Leu Ala Met Asp Glu Gly Gln Tyr Asp Gly
195 200 205

Lys Val Asp Val Trp Ser Leu Gly Ile Thr Cys Ile Glu Leu Ala Glu
210 215 220

Arg Lys Pro Pro Leu Phe Asn Met Asn Ala Met Ser Ala Leu Tyr His
225 230 235 240

Ile Ala Gln Asn Glu Ser Pro Thr Leu Gln Ser Asn Glu Trp Ser Asp
245 250 255

Tyr Phe Arg Asn Phe Val Asp Ser Cys Leu Gln Lys Ile Pro Gln Asp
260 265 270

Arg Pro Thr Ser Glu Glu Leu Leu Lys His Ile Phe Val Leu Arg Glu
275 280 285

Arg Pro Glu Thr Val Leu Ile Asp Leu Ile Gln Arg Thr Lys Asp Ala
290 295 300

WO 03/051905

Val Arg Glu Leu Asp Asn Leu Gln Tyr Arg Lys Met Lys Lys Leu Leu
 305 310 315 320

Phe Gln Glu Ala His Asn Gly Pro Ala Val Glu Ala Gln Glu Glu
 325 330 335

Glu Glu Gln Asp His Gly Val Gly Arg Thr Gly Thr Val Asn Ser Val
 340 345 350

Gly Ser Asn Gln Ser Ile Pro Ser Met Ser Ile Ser Ala Ser Ser Gln
 355 360 365

Ser Ser Ser Val Asn Ser Leu Pro Asp Val Ser Asp Asp Lys Ser Glu
 370 375 380

Leu Asp Met Met Glu Gly Asp His Thr Val Met Ser Asn Ser Ser Val
 385 390 395 400

Ile His Leu Lys Pro Glu Glu Glu Asn Tyr Arg Glu Glu Gly Asp Pro
 405 410 415

Arg Thr Arg Ala Ser Asp Pro Gln Ser Pro Pro Gln Val Ser Arg His
 420 425 430

Lys Ser His Tyr Arg Asn Arg Glu His Phe Ala Thr Ile Arg Thr Ala
 435 440 445

Ser Leu Val Thr Arg Gln Met Gln Glu His Glu Gln Asp Ser Glu Leu
 450 455 460

Arg Glu Gln Met Ser Gly Tyr Lys Arg Met Arg Arg Gln His Gln Lys
 465 470 475 480

Gln Leu Met Thr Leu Glu Asn Lys Leu Lys Ala Glu Met Asp Glu His
 485 490 495

Arg Leu Arg Leu Asp Lys Asp Leu Glu Thr Gln Arg Asn Asn Phe Ala
 500 505 510

Ala Glu Met Glu Lys Leu Ile Lys Lys His Gln Ala Ala Met Glu Lys
 515 520 525

Glu Ala Lys Val Met Ser Asn Glu Glu Lys Lys Phe Gln Gln His Ile
 530 535 540

Gln Ala Gln Gln Lys Lys Glu Leu Asn Ser Phe Leu Glu Ser Gln Lys
545 550 555 560

Arg Glu Tyr Lys Leu Arg Lys Glu Gln Leu Lys Glu Glu Leu Asn Glu
565 570 575

Asn Gln Ser Thr Pro Lys Lys Glu Lys Gln Glu Trp Leu Ser Lys Gln
580 585 590

Lys Glu Asn Ile Gln His Phe Gln Ala Glu Glu Glu Ala Asn Leu Leu
595 600 605

Arg Arg Gln Arg Gln Tyr Leu Glu Leu Glu Cys Arg Arg Phe Lys Arg
610 615 620

Arg Met Leu Leu Gly Arg His Asn Leu Glu Gln Asp Leu Val Arg Glu
625 630 635 640

Glu Leu Asn Lys Arg Gln Thr Gln Lys Asp Leu Glu His Ala Met Leu
645 650 655

Leu Arg Gln His Glu Ser Met Gln Glu Leu Glu Phe Arg His Leu Asn
660 665 670

Thr Ile Gln Lys Met Arg Cys Glu Leu Ile Arg Leu Gln His Gln Thr
675 680 685

Glu Leu Thr Asn Gln Leu Glu Tyr Asn Lys Arg Arg Glu Arg Glu Leu
690 695 700

Arg Arg Lys His Val Met Glu Val Arg Gln Gln Pro Lys Ser Leu Lys
705 710 715 720

Ser Lys Glu Leu Gln Ile Lys Lys Gln Phe Gln Asp Thr Cys Lys Ile
725 730 735

Gln Thr Arg Gln Tyr Lys Ala Leu Arg Asn His Leu Leu Glu Thr Thr
740 745 750

Pro Lys Ser Glu His Lys Ala Val Leu Lys Arg Leu Lys Glu Glu Gln
755 760 765

Thr Arg Lys Leu Ala Ile Leu Ala Glu Gln Tyr Asp His Ser Ile Asn
770 775 780

Glu Met Leu Ser Thr Gln Ala Leu Arg Leu Asp Glu Ala Gln Glu Ala
785 790 795 800

Glu Cys Gln Val Leu Lys Met Gln Leu Gln Gln Glu Leu Glu Leu Leu
805 810 815

Asn Ala Tyr Gln Ser Lys Ile Lys Met Gln Ala Glu Ala Gln His Asp
820 825 830

Arg Glu Leu Arg Glu Leu Glu Gln Arg Val Ser Leu Arg Arg Ala Leu
835 840 845

Leu Glu Gln Lys Ile Glu Glu Glu Met Leu Ala Leu Gln Asn Glu Arg
850 855 860

Thr Glu Arg Ile Arg Ser Leu Leu Glu Arg Gln Ala Arg Glu Ile Glu
865 870 875 880

Ala Phe Asp Ser Glu Ser Met Arg Leu Gly Phe Ser Asn Met Val Leu
885 890 895

Ser Asn Leu Ser Pro Glu Ala Phe Ser His Ser Tyr Pro Gly Ala Ser
900 905 910

Gly Trp Ser His Asn Pro Thr Gly Gly Pro Gly Pro His Trp Gly His
915 920 925

Pro Met Gly Gly Pro Pro Gln Ala Trp Gly His Pro Met Gln Gly Gly
930 935 940

Pro Gln Pro Trp Gly His Pro Ser Gly Pro Met Gln Gly Val Pro Arg
945 950 955 960

Gly Ser Ser Met Gly Val Arg Asn Ser Pro Gln Ala Leu Arg Arg Thr
965 970 975

Ala Ser Gly Gly Arg Thr Glu Gln Gly Met Ser Arg Ser Thr Ser Val
980 985 990

Thr Ser Gln Ile Ser Asn Gly Ser His Met Ser Tyr Thr
995 1000 1005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 June 2003 (26.06.2003)

PCT

(10) International Publication Number
WO 03/051905 A3

(51) International Patent Classification⁷: **A61K 38/00**

(21) International Application Number: **PCT/US02/39742**

(22) International Filing Date:
12 December 2002 (12.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/340,312 13 December 2001 (13.12.2001) US

(71) Applicant: **EXELIXIS, INC.** [US/US]; P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(72) Inventors: **COSTA, Michael, A.**; 18 Hazelwood Avenue, San Francisco, CA 94112 (US). **GENDREAU, Steven, Brian**; 2801 Turk Street, #103, San Francisco, CA 94118 (US). **DORA, Emery, G., III**; 1847 32nd Avenue, San Francisco, CA 94122 (US). **NICOLL, Monique**; 224 Naomi Avenue, Pacifica, CA 94044 (US). **URBANI, Lenore**; 59 Playbowl Drive, La Honda, CA 94020 (US). **LARSON, Jeffrey, S.**; 1220 El Camino Real #305, Burlingame, CA 94010 (US).

(74) Agents: **SHAYESTEH, Laleh et al.**; Exelixis, Inc., P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report.

(88) Date of publication of the international search report:
16 October 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **TAOJIKS AS MODIFIERS OF THE BETA-CATENIN PATHWAY AND METHODS OF USE**

(57) Abstract: Human TAOJIK genes are identified as modulators of the beta-catenin pathway, and thus are therapeutic targets for disorders associated with defective beta-catenin function. Methods for identifying modulators of beta-catenin, comprising screening for agents that modulate the activity of TAOJIK are provided.



WO 03/051905 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/39742

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 38/00

US CL : 435/6; 514/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6; 514/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TASSI, E. et al. Human JIK, a Novel Member of the STE20 Kinase Family That Inhibits JNK and Is Negatively Regulated by Epidermal Growth Factor. J. Biol. Chem. 19 November 1999, Vol. 274, No. 47, pages 33287-33295.	1-26

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 April 2003 (28.04.2003)

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Date of mailing of the international search report

30 JUN 2003

Authorized officer

Sheela Huff

Telephone No. 703-308-1235

INTERNATIONAL SEARCH REPORT

PCT/US02/39742

Continuation of B. FIELDS SEARCHED Item 3:

WEST, STN

search terms: beta catenin. kinases

THIS PAGE BLANK (USPTO)